

Field Book

of

W.L. Jepson

1921

If lost, finder please return to
W.L. Jepson, 2712 Berwin Ave., Berkeley,
California and receive reward.

No. 39.

No. 9926

3

is -

In

ways

and

marshes

slopes

this

summer

re

tive

that -

re

with

San Rafael to Willits, Sept. 8, '21.

1
- Left Berkeley 7:30^{a.m.}; his machines
in party: Dr. Mrs. Wiegand of
Cornell; Prof. Upton of Cornell;
Mrs. W's mother; Betty, the
adopted child; my assistants
Miss Newlon and Miss Stuart;
and myself.

No. 9426. *Hernizaria luteaefolia* DC.
This yellow-fl. Tarweed

filled with a yellow sheet the
little valleys of Marin Co. north
of San Rafael, and especially
the sloping valley plate forming
the mouth of cañon.

No. 9427. *Picris echinoides*

By railway track.

No. 9428. *Lessingia hololeuca*
Greene Ignacio

No. 9429. *Montzelia laevicaulis*
T. & G. Russian River bed, Healdsburg

No. 9430. *Trichostema lanceolatum*
Bth. Russian River bed at Healds-
burg.

No. 9431. *Boissduvalia* ^{*densiflora* (Lindl.) S. Wats.} ~~*stricta*~~ <sup>(Sonne-
co.)</sup>
E. of Lytton School ne. of Healdsburg, on
s. side Alexander Valley. Gulch bed.

No. 9432. *Mimouthe pilosa* (Benth.) Greene.
with 9431. Det. A. L. G.

No. 9433. *Gnaphalium*
with 9431.

No. 9433a. *Hemizonia fitchii* Gray var. *ab-*
undata. Lytton School, gulch s. of (Alexander)

No. 9434. *Chrysopsis*
Stream bed, Hopland.

No. 9435. *Brickellia*
with 9434.

No. 9436. *Euphorbia serpyllifolia* Pers.
with 9434.

No. 9437. *Nicotiana bigelovii* Wats.
with 9434.

No. 9438. *Chrysopsis*
with 9434. This is very ^{cont. p. 4}

- Dr. Wiegand speaks of his dis-
appointment in marshes for
botanizing, i.e. here in California. In
his state of New York marshes always
furnish the best botanizing ground.
Marshes in California, freshwater marshes
are likely to be poor in species.
On the contrary rocky places or slopes
are rich in interesting species. This
note has reference chiefly to summer
botanizing in California. at low
altitudes.

- "Any botanist who gives an
intensive stress to nomenclature
seems to me to be rather defective
in true scholarship or somewhat
unbalanced". I made the above re-
mark to Wiegand who agreed with
it.

4 Ukiah to Long Valley.

cont. p. 2.

abundant in the broad flood beds of streams in this region. I saw areas somewhere between Ukiah and Long Valley where the broad flat of the stream bench was covered with it for a mile - possibly for a greater distance.

No. 9439. *Mimatanthe pilosa* (Benth.) Greene.
Det: A. L. G.
with 9434.

No. 9440. *Xanthium canadense* Mill.
with 9434.

No. 9441. *Eriogonum roseum*
first stream bed n. of
Ukiah = Ackermann Creek.

No. 9442. *Centaurea*
with 9441.

No. 9443. *Lagophylla ramosissima*
Nutt. with 9441.

No. 9444. *Chrysopsis*
with 9441.

No. 9445. *Eremocarpus setigerus* Bth.
with 9441.

No. 9446. *Asclepias eriocarpa* Benth.
Long Valley, about 4 mi. n. of

Sept. 9, 1921.

menloville C.

5

Long Valley. Some of these valleys have local names doubtless and this one, where we lunched, may not be a real part of Long Valley, but it is close to and in the same bowl of the mountains as Long Valley.

- Rattlesnake Creek: *Trillium ovatum*, *Trientalis latifolia*

No. 9447. *Eriogonum nudum* var. *nudum*
Long Valley.

No. 9448. *Eriogonum latifolium* Smith
Ackermann Creek, Ukiah Valley.

No. 9449. *Stephanomeria virgata*
with 9448.

No. 9450. *Aralia californica* Wats.
Rattlesnake Creek, about
half-way betw. Cummings and Leg-
get Valley, on mountain side, in

6 Mendocino Co.

South Fork Eel River, 9 Sept. '21.

a lateral shady gulch with water
trickle.No. 9451. *Minulus cardinalis* Dougl.
with 9450.No. 9452. *Disporum hookeri* Britton.
with 9450.

No. 9453. *Penstemon corymbosus* Bth.
 Anthers dehiscent from the base up-
 ward so that the cells are confluent
 and explanate; they are quite glabrous.
 The sterile stamen is densely hairy
 on one side. The upper lip is notched;
 lower lip 3-lobed, the lobes oblong-
 ovate, ventish. Inside surface of
 lower lip, the whole lip, linear-sym-
 mellulate. This pubescence is hard
 to describe, the hairs are short,
 flat, perhaps scurfy - puberulent is
 better. This is very showy and very
 fine. It is now in splendid flower.
 The woody base is very short, 3 or
 4 in. high. - Red Mt. Creek.

No. 9454. *Malacothrix floccifera*
with 9453.Bull Creek, Sept. 10, 1921. 7
Humboldt Co.Bull Creek Flat.*Sequoia sempervirens*.

Circ. at 4 ft. Approx. Height

No. 1. -	12.10	250-275
" 2 -	13.0	"
" 3 -	45.5	"
" 4 -	40.8 at 5 ft	"
" 5 -	35.5 at 6 ft	"
" 6 -	51.3* at 6 ft.	"

This looks more like
 a *Sequoia gigantea* trunk
 than any of the others

" 7. - 41.8 ft at 6 ft.

* = "Big Tree"

- *Taxus brevifolia* Nutt.

No. 1. 4 ft. circ. at 4 ft.

Height = 36 ft.

Highly irregular tops in this region.

No. 2. 3 ft. 4 tenths at 4 ft.
47 ft. h!

- On the flats Vancouveria

No. 9455. *Tiarella unifoliata* Hook.
Bull Creek flat.

No. 9456. *Taxus brevifolia* Nutt.
Bull Creek flat. See above.

No. 9457. *Abies grandis* Lindl.
Bull Creek flat.

Sept. 10, 1921.
Stemholdt Co.

- There are comparatively few natural objects on this earth that one wants to see more than once; there are few that one wants to see a good many times. But one revisits Sequoia gigantea many times and always with a fresh sense of exaltation. Each visit brings fresh ideas to bear on these trees, so very hard is it to encompass their significance by one or even by a few visits. The same view can be taken of the great Redwoods on the flats of Bull Creek - they are true giants and their average runs high. One is strongly impressed by the unusual character of this assemblage.

- "Vaccinium ovalum", 40 ft. in the air on a dead Redwood trunk! This takes to Redwoods that have died, either erect or lying prostrate. Remaining

- *Passania densiflora* that hangs itself. A Tan Oak seedling ^{led} started to grow on a Redwood log, one end of which was a bit elevated as it lay on the forest floor. The Redwood rotted and sloughed away on the outside of the log, destroying in part the foothold of the root-system of the Tan Oak seedling now 4 ft. high. The seedling toppled over and hung upside down in the air, its roots still ramifying in the trunk.

- The slender limbs of the *Sequoia sempervirens*, straight or drooping. The big limbs of the *Sequoia gigantea* and crooked and stout and short. The limbs of *Sequoia sempervirens* are short. The symmetry of Redwood is so great that the trees do not look large. But there is a constant miracle performed before our eyes when one sees a human being appear in the scene. Two or three persons walk along the flat and the eye at once says they are under-sized persons, often ridiculously undersized. You look carefully the second time and you see that they are quite normal sized people. After this is repeated the trees begin gradually to take on their true proportions.

- Blimney still, living foliage
 half gone longitudinally and
 charred bleached from the
 side I viewed it. 80 ft. h.

Another near it, 40 ft. h.
 with holes in it at intervals.

These great standing hulks are
 impressive in an unusual degree.

- their height is great and the
 huge windows in them assist
 the eye to comprehend their size.

- Seedlings. Young seedling tree
 23 ft. h., $1\frac{1}{2}$ in. diam. at base.

A small thicket of seedlings stands
 on the banks of Bull Creek. These
 seedlings, 20-25 feet high, were very
 slender and support or protect each
 other. The wind has released one
 and it has fallen over and
 forms a low curve with the tips
 on the ground. Its measurements
 are as above.

[Bolling Grove.

Redwood opp. memorial tablet,
 on w. side of road, 36.6 ft. at
 6 ft., 175 ft. h.]

- Splitting of Redwood. Redwood is
 remarkably straight-grained. One can
 split it in any degree of thickness -
 railway ties or posts for fencing -
 or to any degree of thinness -
 grape-vine stakes or shingles or
 thin slivers, the grain running true
 from end to end. No other Californian
 wood approaches Redwood in this
 quality; perhaps it is excelled by no
 other wood whatsoever. ^{in this particular}

[The fern collected at Leggett Valley
 see p. 20, no. 9428^a, *Pellaea*
densa Hook., grew on the red
 flat, at the point where one turns
 east from Leggett Valley. Just after one
 goes down grade further; and crosses a bridge.]

- Plants in Bull Flat.
Pancoveria hexandra
Cornus nuttallii Aud.
Vaccinium ovatum Pursh.
Borylus nortata
Abies grandis Lindl

Philipsville and Garberville, Sept. 10.
Humboldt Co.

- No. 9458. *Helianthus*
 Flat at Philipsville, shady.
 No. 9459. *Arctium minus* Bernh.
 This burdock grew with 9458
 No. 9460.
 This fern grew near Garberville.
 No. 9461. *Adiantum jordanii* Muell.
 This fern grew with 9460.
 No. 9462. *Glinetia andrewsiana* Torr.

Sept. 10, 1921.

- There is no disappointment in the Redwoods of Bull Creek. I have said so much of them to others that I am doubtful how they may impress me when I visit them. They give an impression of very great height but mere height is not the strongest impression. One gets an idea of upper worlds, removed from the ground on which I walk, upper zones of life flourishing in their own spheres and representing something different from the world below. I was struck by the mightiness of the upper stories of the forest as a whole - upper regions not visited or explored and representing something entirely new.

Redwood Flat, west bank of S. Fork Eel, roughly
about 7 or 8 mi. S. of Garberville, not far from
Indian Creek. * Humboldt Co.

- The open Redwood forest is as interesting as the denser forest; and there is much half-open Redwood along the South Fork. We break camp at Bull Creek early in the afternoon of Sept. 10 and return southward, passing Miranda, Phil. iperville, Garberville; S. of Garberville crossing the river on a large bridge and stopping over night at a fine Redwood flat.

- Singular burls at Sat. night's camp. The forest on the flat is composed of huge Redwoods. A number have conspicuous burls. On one tree the burls are near the ground, very large and one is shaped like a giant man facing the trunk.

- *Osculus californica* Nutt. occurs in little patches at bases of hills in open or semi-open country.

No. 9463. *Vitis californica* Benth.
River bank

No. 9464. *Artemisia dracunculoides* L.
with 9463

* = Devoy Flat acc. Solon H. Williams in litt.
Humboldt Co.

No. 9465.

Sept. 11, 1921. Devoy Flat.

S. Fork Eel.

17

- *Trichostema laxum* Gray.

Corolla indigo blue. Upper lobes obovate, acute; lateral lobes lower lip broadly oblong, acute; middle lobe lower lip long-oblong, truncate. Its sides obscurely lined up, especially below the middle as as to be weakly trough-shaped, its color very deep purple, the spots and markings on a light ground, the light ground not abundant in area. All lobes of the same length.


Lip with undulate side-margins.
No. 9466. See p. 18.

- *Trichostema lanceolatum* Bth.
See p. 18.


Lobes upper lip oblong narrow, somewhat coiled or rather twisted not really coiled, longer than the lobes of lower lip. All lobes light purple or light blue, very light, the middle lobe of lower lip almost with light purple weak markings

Redwood Flat, not far from Indian Creek.
 = Servey Flat * Humboldt Co.
 18 or the markings below dark purple.
 Lip a little trough-shaped.

The very rough and hasty field sketches show the difference in the curving of the tube in the two species.



Pale-blue fld.
 No. 9466.



Purple-fl.
 No. 9465.

* The creek flows n. end
 of the flat we call
 Grove Creek. - Chas.
 Neilsen in litt. Nov. 22, '21.

- Rocky gulch. Sept. 11, 1921, ^{near Mendocino} Little Red Mt. Creek.
 c. 2000 ft. S. Fork Eel R.
 ✓ No. 9467. *Baccharis pilularis* ssp. *consanguinea* (Dr.) Wolf
 ✓ No. 9468. *Pentstemon corymbosus* Bth.
 ✓ No. 9469. *Brickellia californica*
 ✓ No. 9470. *Chrysopsis*
 ✓ No. 9471. *Ribes* ^{foezlii var.} *amictum* Greene.
 - A. Tex. *no-calyculatum* Kell.
 ✓ No. 9472. *Antirrhinum vagans* Gray.
 ✓ No. 9473. *Lonicera hispidula*
 ✓ No. 9474. *Zauschneria californica* Pursh.

Near Piercy P.O.

* Little Red Mountain Creek = Mendocino Co.
 nw.

- Great colonies of White Thorn
 in the valley n. of Leggett
 Valley (*Crataegus incanus* T. & G.).
- Private herbaria. Wiegand says he
 bought from Dr. J. K. Small of the U. S.
 Bot. Gard., for Cornell, Small's private
 collection of mosses.
- Leggett Valley, nw. Mendocino Co.
Libocedrus decurrens - 90 ft. h.
Pseudotsuga taxifolia
Abies macrophyllum Pursh
Heteromela arbutifolia Roem.
Umbellularia californica Nutt.

Leggett Valley, S. Fork Eel River.
nv. Mendocino Co.

- *Silene californica*, Tap root 2 ft. long, a little larger than a pencil but holding its size downward.

No. 9475. *Hemizonia*

Dry open land.

[Barbain & Darlington. On a trip to Labrador, Gray Herbarium party, Dr. Wiegand says there was a Barbain and a Darlington in the party, descendants of the famous botanists of those names.]

No. 9476. *Polygonum*

with 9475

No. 9477. *Eriogonum vimineum*

Sliding loose soil.

No. 9478. *Rhamnus californica* Esch.

No. 9478a. *Pellaea densa* Hook. ^{Typica as Wolf} _{See p. 13.}

- The road from Leggett Valley east to Cummings is well up on the mountain side. It is narrow and with short turns - but a smooth bed.

Sept. 11, 1921.

- F. V. Coville. This man is very arbitrary and dogmatic, acc. to Dr. Wiegand, thus confirming the impressions in general that people give of him.

- We drove rapidly from Cummings south over the little divides to Long Valley, thence south to Willits, Ukiah and Hopland. Some 20 minutes drive this side ^(northerly) of Cloverdale I opened the gate of the Alder Glen School yard, drove to the rear and we camped for the night. As I was laying out my sleeping bag I heard a great racket on the road above, and looked in time to see a Ford machine go off the highway and turn over twice. While the occupants scuttled around on the road above, the Ford engine went right on, running as tho nothing had happened.

Alexander Valley, south side.

- No. 9479. *Plantago lanceolata* L.
moist flat.
- No. 9480. *Artemisia* ^{*heterophylla* Nutt.}
~~*douglasiana* Bess.~~ Moist flat.
- No. 9481. *Baccharis Douglasii* DC.
Moist flat.
- No. 9482. *Xanthium Canadense*
Moist flat.
- No. 9483. *Psoralea macrostachya* DC.
Moist flat.
- No. 9484. ~~*Solidago*~~ *occidentalis* Nutt.
Moist flat.
- No. 9485. *Mentha arvensis* L. var. *villosa*
~~*densa* Briq.~~ Moist flat.
- No. 9486. *Verbena* ^{*lasiostachys* var.}
~~*septentrionalis* Moldenke~~ Moist flat.
- No. 9487. ^{*A. fascicularis* Dcne.} *Asclepias mexicana* Cav.
Moist flat.
- No. 9488. *Cornus glabrata* Benth.
8 ft. h. Moist flat.
- No. 9489. *Hemizonia clevelandii* Greene
Hayground but
not as dry as usual.

- alt. c. 150 ft. Sept. 12, 1921. Sonoma Co.

- We stopped to botanize in Alexander Valley on a low flat surrounded by trees. The area had been mown for hay but it was a rather moist place and offered good botanizing for the short time we were there. At a hamlet near this place, the roads fork, one fork going to Kaldsburg, the other easterly and on thru Knight's Valley to the Napa Valley. We follow the latter road and travel down the ever-beautiful Napa Valley, stopping for a luncheon in the open air, as usual, ^{this time} in Hunt's Geor., St. Helena. Thence on to Berkeley, where we arrived about 3:30. In the evening Dr. Wiegand lectured before the California Botanical Society on "A Botanical Expedition across the Continent".

24 Fairfax Manor to San Anselmo Creek

- ✓ No. 9490. *Corum Kelloggii* Gray.
Hill slope sw. of Fairfax Manor.
- ✓ No. 9491. *Asteriscia californica* Less.
West of Fairfax, circa 800 ft.
- ✓ No. 9492. *Heteromeles arbutifolia* Roem.
With 9491.
- ✓ No. 9493. *Aster*
Tributary of San Anselmo Creek,
running north
- ✓ No. 9494. *Quercus garryana* Dougl.
With 9493.
- ✓ No. 9495. *Quercus garryana* Dougl.
With 9493.
- ✓ No. 9496. *Eremocarpus setigerus* B. & P.
Open flat, upper Lagunitas Creek,
near the old San Tempe ranch.
- ✓ No. 9497. *Juncus*
Lower end of flat. See 9496.
- ✓ No. 9498. *Corum gardneri* Gray.
Lake Lagunitas
- ✓ No. 9499. *Juncus xiphioides* Mey.
Phoenix Lake, Marin Co.

and Phoenix Lake, Marin Co., Sept. 24

1921.

25

- ✓ No. 9500. *Cyperus*
Phoenix Lake, Mt. Tamalpais.
- Went by train to Fairfax Manor, thence
up stream and up hill westerly, on
a little south, thence to the left of
a cow-ranch, taking down the
headwaters of a dry stream that
led us into a deep cañon and
brought us out in the "wash" of
San Anselmo Creek; thence straight
up the hill to the second road,
then ^{south} over the ridge to open country
forming a broad low valley,
the lower part of which runs
Lagunitas Creek (probable site of
old San Tempe Ranch? - or this
may have been 400 yds. up in
the trees in the hills), thence
to Lake Lagunitas and Ross
+ Pt. San Quentin. - Miss Newlon,
Miss Stewart, Miss Spalding, myself.

no. 9501. *Cornus* ^{*palustris* var.} *californica* Greene
 Stamens 5, alt. with petals; ^{near}
 opp. petals are 5 fan-shaped
 on broadly oblong dilated
 structures fimbriately cleft at
 summit with a yellow gland
 tipped on each. Stigmas 5, very
 short, no style. - Near Rattlesnake
 Camp.

No. 9502. *Arctostaphylos sensitiva* Jepson
 Crown sprouts on fire trail, showing large serr.
 leaves. Collected on the zig-zag trail.

- *Ericameria arborescens* favors
 the hottest sw. chamise slopes.
 It seems abundant in a ribbon
 on the lower side of each trail,
 i.e. on broken ground.*

- A hill slope of *Adenostoma*
fasciculatum now looks dead
 - so red brown is it. This red-
 brown is given by the dried pan-
 icles.

*The shrubs are locally called Golden Fleeces

- *Umbellularia californica* is a most
 versatile tree. Its capacity to occupy
 very unlike situations and assume
 very different ^{at point} is well illustrated on
 Mt. Tamalpais. forms

- *Baccharis pilularis*, occurs prostrate
 about 300 yards westerly from be-
 ginning of pipe-line trail at
 Muir Woods ridge ^{at point} where railway
 crosses ridge.

(No. 9504)

- *Arctostaphylos montana* Eastw.
 Densely branched in top, forming a
 thick top, 3 to 4 ft. h. and twice
 as broad. One thicket of it, very
 close evenly branched in top,
 6 to 6 1/2 ft. h., 24 ft. broad,
 all from one central root. Cont. p. 28.

- *A. virgata* Eastw. Very glandular,
 6 to 8 ft. h., apparently not crown
 sprouting.

Cont. from p. 27.

- *Arctostaphylos montana* Eastw.

This is a very remarkable individual. It is on left-hand side of trail about $\frac{1}{2}$ mile easterly from Amphitheatre on trail to West Peak.

The original individual, about 6 $\frac{1}{2}$ ft. high and erect seems later* to have developed the power to throw out long lateral branches. These extend out a long way and are closely branched with erect branches - the top being thick and sloping down from the open central parent part of the bush. No. 9504. Cont. p. 30.

* It evidently came to its height maturity and then under some stimulus it threw out ^{strong stems} laterally from its trunk base as indicated. The "original" bush is like an open *a. manzanita* bush 6 ft. high. The lateral development forms a low sloping crown which is very thick and quite even.

Oct. 2, 1921.

No. 9503. *Quercus wislizenii* A.D.C. var. *fulvescens*. Ridge betw. Mill Valley and Muir Woods.

No. 9504. *Arctostaphylos montana* Eastw. Bill Williams trail. See p. 27.

No. 9505.

No. 9506. *Quercus wislizenii* A.D.C. var. *fulvescens* Engelm. Bill Williams trail. It is common along this ridge between West Peak and Rock Spring. It is in good fruiting and shows as great a range of variation in shape and size of nuts as the species, varying from small cylindric nuts to thick or globose ones.

No. 9507. *Calycadenia multiglandulosa* DC. ssp. *cephalotes* (DC.) Keck
Bootjack Camp.

No. 9508. *Zauschneria californica*
Presl. Bootjack Camp.

No. 9509. *Calycadenia multiglandulosa*
DC. ssp. *cephalotes* (DC.) Keck.
Mountain Theater.

No. 9510. *Epilobium franciscanum*
Barbey. Bootjack Camp.

No. 9511. *Grindelia*
Bootjack Camp.

No. 9512. *Cypripedium argenteum* Jeps.
Near West Peak.

- *Arctostaphylos montana* Eastw.
Cont. from p. 28. Several more very
large individuals were seen along
the Bill Williams Trail on the way
to West Peak from Mountain Theater.
Perhaps they were equally large
but all had the same habit;
elongated lateral branches mak-
ing a low broad crown, very
broad and relatively very low.

No. 9513. *Arctostaphylos*

I thought in the field this
may be *A. virgata* Eastw. But there
seems to be every intergrade betw.
this and *A. glandulosa* which
is fully developed along the
trail to West Peak. No. 9513 was
obtained nearer the Mountain
Theater end of the trail.

- J. G. Lemmon was a most pushing
person, ^{albeit of modest manner, still} very aggressive in getting
the name of Lemmon to the front.
But he was utterly imperious and
had to resort to sly tricks. I carried
him free on the *Erythraea* list
but when the first edition of the
Flora of Western Middle California
came out I neglected to send him
a copy. There was no reason why
I should ^{give him one}. I had to give away a
hundred copies. That was enough.
Lemmon could have bought it in

any of the book-stores. But here he came in to my lab. one day, plainly very much agitated. So he wanted to buy a copy. No, he would not accept a free copy. What had he done to be treated in that way - that no copy had been sent him at once? No, he could not and would not accept a copy. No, he must insist on buying a copy. However I finally quieted him and he went away happy and child-like with his presentation copy. He had no part in the book in any way, but he felt entitled to a copy. He was the Emperor Norton of California Botany. Eaton, *Trans N. Am. 2:125* speaks of Lemmon's modesty. He was extremely modest in manner but it was a smoke-screen to conceal his push and imperiousness.

- Professor Arthur^{de} Jazgowski, director of the Institute of Mycology and Pathology, Petrograd, is here. He is to speak before the meeting of the California Botanical Soc. Sat. eve. A snip-nosed clerk in the President's Office rebuffed him because his name ended in "-ski". He is not however a Bolshevik. Although internationally known as a scientist he is said to be a prince of the old empire and he certainly had a half-million acres confiscated by the Soviets. He pointed out the folly of fiat money and indicated that paper, in Russia was worth more just as paper than the current value of the denominations as money! That is the notes when printed would buy less in the open market than the paper cost to print as money!! He is a very human and likeable gentle-

man and his itinerary is being arranged by the Federal Quarantine Board. He is being accompanied in his tour by high officials of the Board and of the Bureau of Plant Industry. See my letters to the President's office dated Oct. 6 and 7. Continued at bottom of p. 41 seq.

- David T. Mason, formerly Professor of Forestry, lunched with me today. He said that on Big River, Mendocino Co., an acre plot of 2nd growth Redwood 48 years old averaged 150 ft. in height, some of it being 160 ft. h. The amount of B.M. was found in excess of that obtained per acre, as an average, from lumbering mature stands. Frith & Metcalf's measurements. Mason is now a forest engineer and has contracts with several Redwood companies to put the operation of their holdings on a permanent

basis. The amount of standing Redwood timber exclusive of associated species is 70 billion feet. Mason regards this figure as too low; but at present rate of cutting the supply would last 100 years - before all virgin Redwood would be milled. Years ago the estimates were very low - 25 billion - but the estimates have been constantly raised. Rate of milling is of course likely to increase. - Oct. 25, 1921.

- *Heteromeles arbutifolia* Roem. grow 15 ft. above tide-level, Pobreño Hills, Richmond, near the present San Rafael Ferry Bldg., on the Blake Bros. property, i.e. on the steep bluffs facing the Golden Gate. The shrubs were large once, fruited finely but are now destroyed thru the ravage of the beaver hunter.
- Nov. 21, 1921. Facts from Anson Blake.
- Rattan, V., was at Three Creeks and Redwood Creek, June 27, 1883. See label *acris-galun* spm.

- Austin, R. M., Mrs., noticed in Eaton, Ferns N. Am. 1:74.
- Orcutt, C. R. For latest Orcuttiana see letter file 1922, with comment by S. B. Parish.
- See footnote to *Streptanthus glandulosus* Hook. in Syn. Fl. p. 171, as a good example of the way in which the Gray Herb. people differ from others about species. It is done in a truly dignified way; so different from Greene's way - the more the pity. It is too bad his real abilities were disfigured by a personal way of expressing opinion. Nevertheless Greene's criticisms were often very able and often exceedingly amusing. He enjoyed thoroughly writing them. He had a keen sense of humor and enjoyed the humor in his name being linked with that of Orcutt; for Greene had a high opinion of his abilities and scholarly (cont. p. 41)

- *Leucollaea Davisiae*. J. P. Young, on Whiskey Butte, 4 mi. westerly from ¹³Elva Mills. He found it in flower!
Two trees of *Picea Brewsteriana* on Whiskey Butte. - John P. Young, Oct. 9, 1921. He also says: "J. B. Parry is working on a Flora of South Africa. He is now and has been for a year at Cambridge at the School of Botany & Eng., where he has room and funds, it very congenial, going down to New for a fortnight or so at need. He plans to go back to South Africa via California in about a year."
- Parry, Dr. C. C., notice of by J. M. Coulter, Bot. Gaz. 15:66.
- Jones, W. E., collected in Cal. in 1882, was in San Luis Obispo Co., 5 May. (coll. type *Penstemon californicus* C. & R. Bot. Gaz. 13:143 (1888).
- Lynn W. S. On letter head "State Board of Forestry" his name appears printed as "Forester". See letter in pocket, *Julium Catalinense* sheet in U. C. herb.

— Called at the Parish's last night. Mr. Parish says "Santa Ana Hills" of his labels are south of the San Bernardino Valley, just across the Santa Ana River. These hills have local names, at least in part, which appear on maps. Parish's term does not appear on maps. Miss Powell, the botanical artist, was there. She told this story. Hall left a case unlocked. Miss Walker discovered it and said, "I must scold him about that". Miss P. told H. what he was in for. So when H. & Miss P. went out to the herb. cases next time, Miss Walker was seen by Hall to approach and before Miss H. had a chance to speak H. said: "Miss W. what do you mean by leaving that herb. case unlocked? I am ashamed of you!" Miss W. stopped in stricken astonishment. Surprise and resentment were written large on her countenance!

Nov. 24, 1921.

Why, Professor Hall, she said, you left that case open yourself; you know you did! But H. brushed all this away or rather ignored it utterly and said: Miss W. I am surprised that you should neglect your duty so. Please do not do it again! This story has more point than Miss Powell realized. Everyone who has visited the herb. has been surprised or annoyed to find such a policy in force as the close locking of the cases. This policy is Hall's policy. Miss Harriet Walker is the herbarium attendant or helper. She has been so long in the place and has been so impressed ^{* = see p. 41.} with the necessity of keeping everything locked that she was naturally shocked. Hall, however, regarded it as a good joke.

— Rattan, V. On the labels in Stanford Herb. one finds: The Rattan Herbarium. Presented by Volney Rattan in 1903. On the label of *Pellaea mucronata* is the locality: Rattan's Ranch, Cala-

Veras Co., Mar. 1883." I suspect that Bolander turned over much of his herbarium to Rattan for one of Bolander's herb. labels. "Herbarium, H.M. Bolander" appears in the Rattan collection in the Dudley Herb. See *Pellaea andromedae-folia*.

- Lemmon, J.G. His labels, printed as otherwise, or often atrociously annoying. For example: *Pellaea andromedae-folia* Fr. Crafton & Santa Barbara and Oakland Hills. Legit J.G. Lemmon. All printed. [Dudley Herbarium.]

- I have borrowed *Pellaea* from the Dudley Herb. The sheets in the mechanical make-up of an herb. do not compare with ours: the paper is thin; the numbering stamp is often slovenly; two different genera of ferns are often mounted on the same sheet, even when totally unlike in size and

appearance. - 6 Dec. 1921.

*Cont. from middle p. 39. It has been put upon me for fifteen years that such ^{an act} was an unpardonable sin which no penance or repentance could assuage!

-(Cont. from p. 36.) standing. That the chances of nomenclatorial naming should marry his name with that of Orent, a rather cheap huckster of botanical wares and something of a mountebank, struck him as rather humorous. He really had a fine sense of humor. So did Mrs. B; but hers was sardonic and heavy. Greene's was light and fanciful.

-Cont. from p. 34 near top. He is a tall good-looking man, of ease of manner and of comely address, a most excellent and ready linguist - all this showing his breeding and lineage. A cultivated and highly educated and trained scientist.

- No. 9514. *Ribes* ^{malvaceum} Sm. Half-way up to summit, bare southerly slope, road from ^{mt. D.} Country Club. in fl.
- No. 9515. *Zauschneria californica* Benth. Southerly slope in open woods, half-way up. in fl.
- No. 9516. *Heteromeles arbutifolia* Pop. in fr. = *Photinia arbutifolia* Lindl.
- No. 9517. *Salvia melifera* Greene. Common south slope at the lower altitudes above the foothills.
- No. 9518. *Dicentra chrysantha* H. & A. Higher open slopes; it loves broken ground!
- No. 9519. *Salvia columbariae* South slope, Charnise ridge.
- No. 9520. *Oenothera micrantha* Horn. ^{seed} ~~or~~ *Oenothera verticillata* Greene ~~capsules & seed~~ ^{var. *sonnei* Jepson.}
- No. 9521. *Camissonia intermedia* Lam. seed of this crucifer sent to garden, Dec. 1921.

- No. 9522. *Sphaeola calycina* = *Lepeckinia* South slope, Charnise ridge.
- No. 9523. *Delphinium hesperium* Gray. ^{det. R. S. 10/54} Dead ripe capsules.
- No. 9524. *Entillaria* Dead ripe capsules.
- Drove to Diablo with my assistants Miss Newlin and Miss Stewart. We saw Blue Jays, on the way, burying acorns. From the summit of the mountain the whole Coast Range and Great Valley country was buried or drowned in an inky lake - the heavy pall of smoke that was like a fluid body with a level surface. No Coast Range peaks appeared above this lake floor; but St. Helena's sloping summit - by a darker line betrayed its position at the very level of the smoke pall. The Sierras immedi-

sky eastward showed fairly well, and we probably saw the John group, - Pyramid Peak and others. It was a strange sight. Toward Inlet Cañon the smoke was less dense and gave an artistic effect to the cañons.

Lake Alpine, Marin Co.

Dec. 11, 1921.

- Walked from Ross to Lake Alpine and back to Fairfax with Prof. Walter Mulford.

No. 9525. *Pellaea arnithopus* Hook.

Gravelly outcrop of ledge, rigidly erect. Stipes very stiff + wing.

No. 9526. *Dryopteris arguta* (Kaulf.) Watt
(Det. Tryon, 1958)

Shady bank

= *P. mucronata* var. *mucronata*

No. 9527. *Pellaea arnithopus* Hook.

Rocky outcrop.

No. 9528. *Polypodium californicum* Kaulf.
var. *Kaulfussii* Fort Barry. Breccias of rocks
Cat.

and mossy banks.

No. 9529. *Rumex* (Discarded)

Leaves fleshy, bright green, somewhat shiny, truncatish at base but almost-subcordate in appearance.

Ft. Barry

No. 9530. *Lemna*

Ft. Barry.

No. 9531. ~~*Schmoegetis*~~ ^{*marah*} *jabacea* (Nutt.)
Ft. Barry.

No. 9532. *Mesembryanthemum aequi-*
laterale. Ovary cells 8, 10,

10. Petals linear-oblancoate

Ft. Barry.

— Walked from Sausalito to Fort Barry, at the outer entrance of the Golden Gate, with the Cal. Bot. Soc. Three High School lads, Harland and Francis Fredericks and another boy were along. They added much to the liveliness of the day. They played too near the surf and Harland got ducked. He took off his shoes to dry them by our beach fire and one

Marin Co., Jan. 25, 1922.

shoe was badly burned. Indeed both shoes were in great danger of burning up. But he managed to tie the shoe ^{around} onto his foot. On the way back we came thru the tunnel. The boys hid in the dark recesses and when I came along rushed out as brigands with great shouts: I replied in kind: "Lay on, foul fiends, and damned be he that first cries, hold enough!" This delighted them greatly.

The little groves of Eucalypti in cove facing the Golden Gate are badly burned by frost on the easterly sides. The trees facing the first impact of the recent killing north winds were whitened by the frost. This is the first time I ever saw mature Eucalypti killed by frost in central California.

Stockton, Feb. 3, 1922.

- Went to Stockton to talk to the Philomathean Club in behalf of the Save the Redwoods League. The day was hazy and cloudy but the ranges still showed the fall of snow of last Sunday. The inner Coast Range, as we viewed it from the plains of the San Joaquin, showed heavy snow. The higher mts. were streaked and patchy because of trees and shrubs, but the middle grasslands were one unbroken unspotted field of white. As the range stretched southward this middle zone of spotless white ran lower and lower until finally in the far distance it touched the plain or floor of the San Joaquin. Now it did not really do so; but the curvature of the earth's surface made the illusion and really shut out the dark low foothills so that the eye saw ^{just} over the curvature, the base of the middle snow zone.

- It has been very cold during Dec. and especially January, with white frosts twice the normal, continuing into Feb. High winds in Dec. & Jan, doing much damage. Sun at end of Jan. Precip. half normal in Jan. It is now, Feb. 22, getting a little warmer, but precip. has been heavy in Feb. Sun drifted 2 to 3 ft. deep in Berkeley Hills in Jan. storm.

- N. J. Andersson, coll. at Coloma in 1852(?). Cf. Mem. Torr. Club, 1:20
- Bridges, Thos. Pellaea bridgesii, named for him by Hooker, will keep his name in remembrance for long in California.

Lake San Andreas.

- Scoliofus bigelovii Torr. Slugs are eating the flowers.

No. 9533. *Calochortus albus* Dougl.
Millbrae. Leaves 27 in. long, 1 1/2 in. wide!

No. 9534. *Cytisus scoparius* Link.
This is truly spontaneous and has spread along the roadway for a mile or more, about the summit between Millbrae and Lake San Andreas. The thickets are dense but do not spread far into pasture land.

No. 9535. *Sanicula arctopoides* H. & A.
Lake San Andreas, San Mateo Co.

No. 9536. *Dirca occidentalis* Gray.
This is very attractive at this time and gives a lacy effect to the winter woods. Its flowers are small but borne in great numbers. We saw some very attractive shrubs. Lake San Andreas. One shrub was full 11 ft. high!

No. 9537.

Lichen on board fence. Lake

San Andreas.

- No. 9538. *Aspidium rigidum* var. *argutum* Eat. Abundant in the woods, plainly evergreen, Lake San Andreas.

No. 9539. *Corallorhiza striata* Lindl.
Lake San Andreas. Woods.

No. 9540. *Ribes californicum* H. & A.
Lake San Andreas. 4 to 6 ft. h.

Ovary glabrous and smooth.

Petals white, truncate, involute

Calyx a little pinkish in bud, mostly greenish in anthesis, but often pinkish. Leaves not glandular, very small for *R. californicum*. Style glabrous.

- *Dipsacus fullonum*. Common about Millbrae.

No. 9541. *Umbellularia californica* Nutt. We stopped to rest under a very large tree whose crown dropped to the ground all around

It grew on the flat, west side of abutment about 2 miles south of the Lake San Andreas dam. Its crown was about 90 ft. broad; height about 78 ft.

No. 9542. *Osmaronia cerasiformis* Nutt. ♂ only collected, the ♀ not noted.

No. 9543. *Ribes californicum* H. & A.
Ovary with long slender non-glandular bristles.

- *Holodiscus discolor* 11 ft. h.

- *Aesculus californica* Shach.

- *Ceanothus thyrsiflorus* Esch. 12 ft. h.
crown-sprouting.

No. 9544. *Pedicularis densiflora* Bth.
Bare slope of hill just above (East) a clump of oaks, on Buriburi Ridge.

No. 9545. *Fragaria californica* C. & S.

No. 9556. *Dodecatheon hendersonii* Gray var. *Bernalianum* Greene. Leaves yellowish a little, and

flesher than in the ordinary form. Buriburi Ridge.

Went to Lake San Andreas with the Cal. Bot. Soc., walking from Millbrae and returning over Buriburi Ridge to Burlingame.

— East Hall is being moved from its old-time position to a place south of Strawberry Creek near the Museum of Vertebrate Zoology, via the College av. bridge. There was much tree-cutting to make passage for the building. Trees were cut as follows:

East Drive, 4 oaks.

College av., nine large Elms and maples on west side of street.

Women's Basketball Court, 16 acacias + allied trees.


Physiology Building, 3 acacias on south side.

The total is 32 trees. The street trees were about 30 years old, maybe forty.

— Drove to Tracy Apr. 1 in order to lecture on behalf of Redwoods League for Tracy Women's Club, at 8 P. When I arrived no one there except 2 men. Gradually hall filled up and I was introduced by the Madam President at 9 P. Slides well projected. Afterward they had games, then a banquet. Each man was called upon to respond to a toast. Three-fourths chose to speak about trees. Now these men looked like and were far the average run of men in a small town, shop-keepers, salesmen, a few farmers, ^{many} all were visibly embarrassed when they were called upon but each in his own way expressed an original sentiment about a tree or some experience in the wild woods, the language was not polished but it was simple and direct and often Lincolnian in its force and simplicity.

- No. 9557. *Plantago*
Gravelly flat, mouth of Corral Hollow.
- No. 9558. *Gilia tricolor* Benth.
with 9557.
- No. 9559. ^(Claytonia) ~~Phlox~~ *perfoliata* Donn ex Willd.
Lower part of Corral Hollow.
- No. 9560. *Oenothera graciliflora* H. & A.
Gravelly flat
- No. 9561. *Plagiobothrys canescens* Bth.
Gravelly flat.
- No. 9562. ~~*Madia*~~ *radiata* Kell.
~~*Layia claysana* (non writer)~~
Gray. Clay hill top. det RCB. 12/57
- No. 9563.
- No. 9564. *Trifolium gracilentum* F. & G.
- No. 9565. *Phacelia ciliata* Benth.
Clay hill top; with 9562. Corolla
blue. Scales at base of fil.
orbicular in general outline. See
sketch with spurs of No. 9566.

- No. 9566. *Phacelia douglasii* (Bth.) Torr.
Steep rocky slope. Scales long
and narrow, widest at middle, straight,
longitudinally attached, the pairs
parallel. See sketch with spurs
of 9565. See also 9593.
= *P. fremontii* Torr!! not *P. douglasii*.
- No. 9567. *Tropidocarpum gracile* Hook.
Flats along the creek.
- No. 9568. *Layia platyglossa*
Clay slope, gentle hill.
Rays 6 to 8 lines long. Rays 8, 8, 8, 7, 8.
Rays gold-yellow, white in upper third.
Pappus-bristles many, not hairy or plumose.
Achenes silky-pubescent. Ray achenes ob-
compressed (dorsally flattened), glabrous, no
No. 9569. *Ellisia membranacea* [Pappus].
Bth. This is the same as no. 9585
but was farther down the cañon.
↑ collected
- Corral Hollow, west half in Alameda
Co., east half in San Joaquin Co.
All my plants were collected in San
Joaquin Co., on this trip.

No. 9570. *Baria*Sandy flats. This was common,
especially thru all the lower
part of the hollow.No. 9571. *Dichophragma parviflorum* T. & G.
~~*Sellina affinis* Bol.~~Petals white, deeply cleft, turning
pink in the press.No. 9572. *Dichelostemma multiflorum*
~~*Dracopis capitata* (Benth.)~~ Heller
Sandy hillside.No. 9573. *Tropidocarpum gracile*
Hook.No. 9574. *Platystemon californicus*
Benth. var.Petals sulphur-yellow with a
band of white on each side
toward the base.

Sandy flats.

No. 9575. *Plagiobothrys tenellus* Gray.
Gravelly flats.No. 9576. *Oenothera graciliflora* H. & A.
Drawing from this. see p. 62No. 9577. *Chenopodium californicum*
Muhl.No. 9578. *Salvia carduacea* Benth.
Called Prince of Persia here.No. 9579. *Capsella bursa-pastoris* (L.)
Moench. Abundant, forming large colonies
in rich spots on the flats.No. 9580. *Astragalus nigrescens* Nutt.
Dry flat. = *A. gambellianus* Sheld.No. 9581. *Trifolium tridentatum* Lindl.No. 9582. *Gilia* ^{capitata. Sims ssp. *staminea* Greene, A. N. S. Grant} ~~*schottii* folia Benth.~~
Sandy flat. = *G. staminea* Greene.No. 9583. *Phacelia curvipes* Torr.This is a diminutive
species. It grows in a sandy
pocket between the rocks at
the very summit or eastern

w. San Joaquin Co. (see bottom p. 55) c. 600 ft.

end of the Castle Rock. Some other dwarfs grew with it. A small-flowered *Heuchera* and 9584. Seed of the *Heuchera* taken, only.

No. 9584. *Athyrium pusillum* Greene

No. 9585. *Ellisia membranacea* Bth.

Top of cliff-talus, Castle Rock (see p. 61).

This is exactly like the material used for drawing *Ellisia membranacea* Bth., coll. Berris Hill, Harriet Kelly.

It has the oblong or exclamation point lavender spot on the "petals" below middle; ciliated calyx-lobes; hairy ovary; "scales" consisting of pairs of long-oblong papillae without hairs. They are really like glands. I am glad to see this because it indicates a wide distribution of a definite type. It also indicates that Mrs. K's latest sending, differing only in rounded papillae, belongs to this type. See p. 63.

April 2, 1922.

San Joaquin Co.

No. 9586. *Lepidium nitidum* Nutt.

Castle Rock, Coral Hollow; see p. 61 at bottom.

No. 9587. *Gilia tricolor* Bth.

Gentle slopes near Castle Rock.

This slope was blue with it - a large area - a delicate blue from a distance as if the sky had fallen on the hill in a patch.

No. 9588. *Urtica urens* L. Benth.

sepal 5 times as large as the inner, the inner ones scarcely ciliate or not at all so. Castle Rock.

No. 9589. *Orthocarpus purpureus* Benth. Forming patches of several acres in the fields of grain. (Drawing from this). Tracy.

No. 9590.

Near Castle Rock, lower part of Coral Hollow.

No. 9591. *Stellaria nitens* Nutt.

Grainfields near Tracy.

No. 9592. *Pectocarya penicillata* A. DC.

Summit Castle Rock.

No. 9593. *Phacelia ciliata* Benth.

Grainfields near Tracy.

In every way except habit this seems the same as no. 9565. It has the same hairy rays seated on a small glabrous disk. The scales are fundamentally the same. The corolla of 9565 seemed a deeper blue in the field; but there is no dif. side by side in the lab. style is blue in both and parted the same way. Scales rounded, entire; in 9565, obscurely ^{forked(?)} _{Cont. p. 61.}

No. 9594. *Amsinckia intermedia*

F. & M. This is known as "fireweed" in the fields about Tracy; also as "Siccato Gorda", perhaps Zaccato Gorda?; valued as a dry feed.


No. 9595. *Monolopia major* DC.

Called Downy-back Daisy at Tracy. Very abundant on hillslopes and in grainfields. The low hills west of Tracy had great sheets of it for four miles. Rays 8, 8, 8, 8, 8, 9, 8
Rays 5 to 7 lines long. Corolla

tube with scattered gland-tipped hairs, both ray and disk. Pappus none.
Drawing from this. _{Cont. p. 62.}

- Cont. no. 9562. Corolla tube with gland-tipped hairs. Pappus none!

Ray achenes rounded on the back, a little obcompressed - that is flattened from the sides, perfectly glabrous, as also the disk achenes.

Ray⁺ achene.

disk achene not apiculate.
Ray achene apiculate!
the apiculation set out on the edge of ventral angle, so to speak, a little.

- Cont. no. 9593. There does not appear to be any difference in the scales of the two forms; or, if any, the difference is negligible.

- Castle Rock is in lower part of Corral Hollow.

- Cont. no. 9562. Rays 6 lines long, $4\frac{1}{2}$ lines broad, 3-toothed. Corolla limb with a fringe of hairs on the back of the lobes, these hairs really very straight and bristle-like!

"No. 9596. *Baeria tinella* Greene

Pappus clear white, with 4 oblong toothed paleae on each side, each ser- looking like one large palea; one bristle on each side. Rays 10, 10.

By far the vast majority of the flowers have 2 pappus-bristles; the paleae mostly 2 on each side. Receptacle strongly conical. Drawing from this. ^{↑ slender}

Rays 8, 8, 10, 8, 10, — $2\frac{2}{3}$ lines long. Rays oblong. — Tracy.

- Cont. no. 9576. Filaments lanceolate, dilated, about $\frac{1}{4}$ or $\frac{1}{5}$ as long as the petals. Stigma capitate, slightly exceeding the anthers. Filiform calyx-tube $1\frac{3}{8}$ in. long. Petals 4 to 5

April 2, 1922

lines long. Stamens very short. Calyx-lobes hairy. Ovary lightly pubescent, 4-angled, the angles projecting into strong horn-like lobes at the summit; or rather, above.

- Cont. no. 9585. Even the slender smaller-leaved forms have the same characters. I found it common up and down Corral Hollow.

Linanthus dichotomus Bth.
"No. 9597. *Jilja dichotoma* Gray.

Gentle open slope. [I brought fresh plants to the laboratory. For five days they opened very promptly at 4 or 4:30 o'clock.] Drawing from there.

- Came to Tracy last night to lecture for the Save the Redwoods League. Today was taken out to Corral Hollow by Mr. and Mrs. A. P. Arnold of Tracy. Cultivated people who have taken the Amalfi Drive and been on the hill.

Berkeley, Apr. 16, 1922.

- No. 9598. *Ribes divaricatum* Dougl.
Plants from 3 to 7 ft. high; stems
mostly bending over
Stamens exceeding the calyx-lobes;
styles hairy; calyx-tube hairy &
little; ovary glabrous and smooth.
April 11, 1922.

- No. 9599. *Collinsia sparsiflora* F. M.
var. collina (Gris.) Newson
San Pablo Valley. Apr. 16

- No. 9600. *Stellaria Cerastium viscosum* L.
San Pablo Valley, Apr. 16.

- No. 9601. *Lepidium nitidum* Nutt.
San Pablo Valley, Apr. 16.

- Cass, E. B., the schoolteacher and friend
of J. G. Lemmon, is living at Downeyville.
He collected *Astragalus casei* and *Coryd.
alis caseana* but stopped short
after the direct personal influence of
Lemmon was no longer exerted.

see p. near

1142.

Peaceful Glen, near Solano Co. 65

200 ft.

Apr. 21, 1922.

- No. 9602. *Minulus tricolor* Hartw.
Mud-dried prods.
- No. 9603. *Thermophilola menziesii* Gray
This plant made a fine show in the
lower parts of grain fields. Scales
oblong, hairy on edge, about parallel,
the tip slightly rounded and non-ciliate.
- No. 9604. *Thysanocarpus radians*
Bth. This is abundant on the
gentle slopes of the low^{est} foothills
and is found in colonies 5 to
20 ft. square, the plants stand-
ing so thickly in these colonies
as to make pinkish or
pale straw-color spots in the
grassy fields. No two colonies
were exactly alike, in a field
of forty acres, differing only
slightly, but I felt that I
could detect some differences
in certain colonies that were obvious

to the eye. This number is pink-winged, the fruit body covered with a fine felt.

No. 9605. *Thysanocarpus radians*

Bth. See no. 9604. Collected in same field. Wings white, that is pale greenish-straw-color, the body finely tomentulose or with a very fine felt as in 9604.

No. 9606. *Thysanocarpus radians*

Bth. Body glabrous, the foot smaller a little than in 9605, otherwise the same. with 9605, i.e. in same field.

No. 9607. *Limnanthes rosea* Hartw.

Mud-dried pool bottom and along its sides. Cor. white, fading rosy.

No. 9608. *Eschscholtzia lobbia* Greene.

Common on gentle slopes.

No. 9609. *Lupinus micranthus*

See p. 114b. Common in spots. Flowers blue, center banner white, fading purple.

No. 9610. *Alopecurus saccatus* Vasey
(DET. Beecher Crampton, Mar. 1922)

This grass grew in the swales.

No. 9611. *Alchornea greenii* Greene.

(Discarded, insect damaged)

No. 9612. *Achyrochaena mollis*

No. 9613. *Trifolium depauperatum* Desv.

No. 9614. *Layia platyglossa* Gray
chrysanthemoides (DC) Gray

Pappus bristles unequal, the two at the longer than the others.
Frequent. [angles of the achene]

No. 9615. *Viola pedunculata* T. & G.

170 ft. Mile Square Hill near Vanden.

No. 9616. *Allium serratum*

170 ft. Ovary lobes lightly creased lengthwise, ending above in 2 erect teeth. Per. $4\frac{3}{4}$ lines long.

Ovary apparently on a short column $\uparrow 4\frac{5}{6}$ on stipe. Mile Square Hill, Vanden, 170 ft.

No. 9617. *Lasthenia conjugens* Greene.

Swirn. 3 m. W. Rays 16, 15. Corolla

68

Swisum, 3 mi. n.e., Apr. 21, 1922.
Solano Co. ^{30 ft.}

tube with short spreading glandular processes, that is hairs, a very few above on the throat. Ovary glabrous. Back of rays with glandular hairs!

Involucral bracts united half-way up! the free portion ovate, very acute.

No. 9618. *Allocarya stipitata* Greene.

Subalkaline flat.

No. 9619. *Astragalus tener* Gray.

Low subalkaline flats 3 mi. n.e.

of Swisum. Grows in patches and is wonderfully clover-like.

Cor. purple, the banner and wings white-tipped. Keel $\frac{1}{2}$ as long as

banner, the wings $\frac{3}{4}$ as long.

Banner notched.

No. 9620. *Lepidium latipes* Hook
with 9619.

Stem erect, the leaves entire, forming a long tuft above the head of pods. Leaves linear. Stem one. with 9619.

75 ft. Rockville, w. Solano Co. 21 Apr. '22. 69

No. 9621. *Trifolium fucatum* Lindl.

With 9619. In this field each clover plant of this species had an individual color pattern on the leaflets. No two were alike!

The patterns consisted of white markings or brownish or blackish markings.

No. 9622. *Iris missouriensis*

Sepals whitish, pale lavender veined, a yellowish spot in middle, ^{the spot} just at the end of a median ridge which arises at base and runs to about middle of sepal and there disappears. Petals 1 in. broad. Stylars ^{branches} white, 4 lines broad, serrate at apex, the slit at apex 5 lines deep. Rockville, on the flat.

No. 9622a. *Pleurapogon californicus* (Nees) Marsh.

No. 9623. *Montia fontana* L.

Low flats where water has stood in winter.

No. 9624. *Orthocarpus faucibarbatius* Gray. Corolla: folds of sacs inside pubescent. This grows in low fields and is very luxuriant. Corolla sacs sulphur-yellow! It is common all thru this district as far as Valleys.

No. 9625. *Scandix pecten-veneris* L.

This umbellifer was weakly represented in the grain fields. Leaves pinnate or bipinnate, the segments (roundish in outline) palmately dissected. Both ♀ + ♂ in same cluster, the latter few or one, the former with stamens also. Outer flowers with the outer petal enlarged! Styles with a very marked dish-like stylopodium!

↑ or dish-shaped

No. 9626. *Thysanocarpus curvipes* Hook.
Peaceful Glen.

The slopes about Peaceful Glen were in radiant bloom. *Eschscholzia lobbia* made yellow spots in the fields. *Nemophila menziesii* has here adapted itself to cultivation. I found it in grain-fields and in vineyards, in fine development! Everything much later than usual.

Mile Square Hill, 1 mi. n. of Vanden, was in glory, with its colonies of *Allium serratum*, more than I have seen elsewhere in like area, and *Viola pedunculata* T. & G.

Even the dreary semi-alkaline flats betw. Vanden and Suisun, dreary in the dull summer time, have now a delicate evanescent beauty. Some fields were widely spread with colors, with white of *Alloxarya stipitata* and yellow of *Isaetheria conjugens*.

No. 9627.

✓ No. 9628. *Alchemilla* ~~*occidentalis* Nutt.~~ *occidentalis* Scop.

Stamen 1. Style filiform, arising from the base. Abundant on the low open hills near Cerrito Creek.

✓ No. 9629. *Evax sparsiflora* Jepson.

♀ flowers without pappus and with ^{slender filiform} corolla, one in the axil of each bract of the involucre; in the central "cup" of the involucre are 4 ♂ flowers with corollas.

I count 6 bracts with ♀ fls, these are imbricated; the receptacle is prolonged upward as a slender or thread-like organ bearing 3 bracts in a whorl at apex, these 3 forming a cup with 2, 3, or 4 ♂ fls. The lower involucre bracts are imbricated, erect, and approximate around the upper bracts in such a way as to form the "cup," that is the lower bracts

are larger and reach up to the apices of the upper bracts. ♂ fls. about $\frac{1}{2}$ line long.

I find that all the lower bracts are fertile, that is bear ♀ fls; the upper bracts appear to be 5, in a whorl. The lower bracts are rigid and form a short-cylindric body bearing in it, on a stipe-like organ (the produced receptacle) and at apex of the stipe-like organ, a cup-like whorl of 5 bracts which have 2, 3, or 4 male flowers. These latter bracts are more woolly. Pappus none. Male fls. red-brown, $\frac{1}{2}$ line long.

- Cont. from p 71. No. 9627. This grass grew in the beds and along the margins, especially the latter, of former spring pools.

✓ No. 9630. *Lonicera involucrata*
Strawberry Canon, Apr. 30.

- No. 9631. *Barberis pinnata* Lag.
2-3 ft. h. Rocky arthropods
in the hills. Sepals 9: 3 small outer
ones, these caducous; 3 larger, ovate, reflexed
from the middle, persistent larger than
the first 3 but soon falling; 3 large,
obovate, curling over the petals, or rather
ascending over the petals. Petals 6, all of
one size. ^{Cont. p. 75}

No. 9632. ^{*Manah*} ~~*Echinocystis*~~ *fabacea* (Naud.)
Greene

No. 9633. *Prunus emarginata* Vesp.

This forms a small colony
dominant on perhaps half an
acre. The plants vary from bushy
shrubs 4 ft. high to slender trees
26 ft. high with trunk five
inches in diameter. I noticed
only one which reached the
largest size named. They are
in full flower now and make
an attractive group at the
northeasterly base of Sugar Loaf.

Apr. 30, 1922

The odor of the "deciduous fruit" or-
chard was distinctly there.

- Cont. from p. 74. No. 9631. All the fls.
examined was full of small whitish
larvae lying between the petals and
the stamens. In Engler & Prankl, the
6 inner perianth organs are regarded
as honey-organs; the outer as
"Ketch-artig", the middle ones as
"Kronen-artig".

I do not see any "basal glands"
on petals ^{of 9633} as described by authors
for the petals of *Barberis*.

- L. H. Wold of the Bureau of Ento-
mology just called. He finds the
same cynipid galls on all
the species of the *Chrysolepis* group
of *Quercus* - *Chrysolepis*, *vaccini-*
folia, *tanacetella*, *vilcoxii* and one
other; and suggests that there
are really all one species. He

Berkeley, May 5, 1922.

investigated the galls of California Oaks during the war as a substitute for the Aleppo Gall which forms the basis for record ink. The per cent of tannin not sufficiently high. Materials high in tannin split with a glassy fracture. The Aleppo Gall does this. A spiny leaf gall on the under side of leaves of *Quercus douglasii* fractures easily and suggests high tannin content. Wald has been traveling by auto all thru the west since last Sept.

— May 5, 1922

- Roehl Cal. collector. *q.*
Bot. Gaz. 48:462.
- Greene, E. L., on his way to Guadalupe and Cerros Islands (West. Am. Sci. 1: 32, — Apr. 1885.).
- Californian Botanical Collectors.
Data may be had of certain men in West Am. Sci., early vols.

North Berkeley Hills*, May 11, 1922.

Contra Costa Co. 7

- ✓ No. 9634. *Ceanothus thyrsiflorus* Esch.
Hollow in the hills.
- ✓ No. 9635. *Psilocarphus tenellus* Nutt.
var. *tenellus*
Leaves of involucre 1 to 2 times as long as head, spatulate-oblong. ♂ fls in center 8, 7, 8. No chaffy bracts at all!
- ✓ No. 9636. *Lotus subpinnatus* Lag.
- ✓ No. 9637. *Plantago erecta* Morris.
Adobe soil.
- ✓ No. 9638. *Sanicula menziesii* H. & A.
- ✓ No. 9639. ^{= *S. californica* Bitter} *Acaena trifida* R. + P.
Rocky outcrop, north slope.
Pinnae segments 7 or 8. I find 3 to 5 stamens (in one infl.); I find no fl. with only 2 stamens.
= *A. pinnatifida* var. *californica* Jepson.

* Contra Costa Co.

78 North Berkeley Hills, Contra Costa Co.

May 15, 1922.

No. 9640. *Mimulus guttatus* DC. Det. A. S. G.
Swamp.

No. 9641. *Phacelia imbricata*? Greene.

Rocky hilltop, dominant on the
grassy top, 100 x 40 ft. = *P. calycosa*
(Gray).

No. 9642. *Wyethia helenioides*, Nutt.

Grassy hillslopes See p. 79.

No. 9643. *Rhus diversiloba* B. + G.

No. 9644. *Malilobus indicus* All.

Moist spots.

No. 9645. *Hypochaeris glabra* L.

Dry hillside

No. 9646. *Wyethia angustifolia* Nutt.

Hill shoulder. See below.

No. 9647. *Matricaria discoidea* DC.

Berkeley. Beaten ground. May 15.

= *M. suavisolens* Buch.

- Cont. 9646. Rays 10, 10, 11, the
ligule $1\frac{3}{4}$ to 2 in. long. Bracts of
the invol. lance-ovate or perhaps
better oblong-ovate, *contracted

North Berkeley Hills.

79

at base to a distinct and definite
but very broad or ligulate "petiole",
this petiolar base purplish-black,
the margin woolly or soft-ciliate or
pilose-white, the upper side pilose,
the lower side glabrous or nearly so.
Herbage in general almost glabrous (or
at most sparingly pubescent), ^{not}
woolly. Achenes with one strong spine-like
bristle. Corolla inflated a little above the base and
at that point bent or knuckled!

- Cont. no. 9642. Herbage and bracts
thinly tomentose, especially the stems
below. Bracts broadly lanceolate from
a broad base, green throughout, no
purplish spot at base! Pappus of
about 4 unequal paleaceous scales.

No. 9648. *Echinocystis foliacea* (Gray)
Greene
May 11.

* or better still: oblong-lanceolate
or oblong

No. 9649. *Heuchera micrantha* Dougl.
Petals white, oblong-ovate,
recurved. Calyx white, its short
teeth incurved and carrying
a green spot, the whole calyx
soon becoming pink. - Face
of rock cliffs. Stamens 5; very half inf.

No. 9650. *Diplocrocus aurantiacus* (Curt.)
Jepson. Very beautiful salmon-color
fls.

No. 9651. *Achillea millefolium* L.

No. 9652. *Microseris linearifolia*
Sch. Bip.

No. 9653. *Arctostaphylos stan-*
fordiana Parry. Cluster of two
shrubs 4 or 5 ft. h., about half-
way up mt. In the field I have
little doubt that this is *A. stan-*
fordiana. Its leaves are re-
markably vertical; they are small.
The small berries, with the very

dark red bluish on one side is also shini-
ly reminiscent of that species. The
leaves are thick - the only point
against the reference that I notice.
There is no crown-sprouting! The
shape of the leaves recalls those of
stanfordiana.

No. 9654. *Helianthella castanea* Greene.
Bracts linear-lanceolate, hispid
above-ciliate*, the tips after anthesis
strongly incurved over the disk.

Bracts of the involucre subtended
by a more or less complete whorl
of very unequal foliaceous bracts.
Rays 11, 12, 14, 16. Scales of the
receptacle with a dense tuft of hairs
at apex, these tufts appearing in the
disk as seen from above, since just
as long as the corollas. They are
* glabrous below and essentially
glabrous above.

units in the tessellation, if one may compare it to a tessellated floor.

The achenes are very much flattened, somewhat cordate at apex. The disk ones are ^{all} fertile except a central cluster of 15 to 30 flowers which are quite ^{about} sterile! The ray flowers are wholly sterile and their fls. have very small ovaries from the start of anthesis.*

Cont. p. 86.

No. 9655. *Cirsium coulteri* Harv. & Gray. This thistle is common along the roadway, upper third of the mt. It loves broken ground and suggests local introduction. It is about 3 ft. high. Both lavender and white heads are.

* This plant grew on the very top of the mt. at the wagon camp and was abundant.

No. 9655a. *Pedicularis densiflora* with 9663.

found in the same 10 sq. ft. of a colony. No. 9656. *Phacelia breweri* Gray.

Land-slips near summit.

Corolla 2 lines long, pale blue. Appendages to stamens a pair of vertical scales, linear in outline, ^{not} hairy. Calyx cleft nearly to the base, its lobes linear, hispido-lanceolate or better hirsutulo-lanceolate. Calyx a little over half length of corolla. Stam. glabrous, a little shorter than the cor. Ovary densely hirsute with white erect hairs. Style very sparingly short-hairy.

No. 9657. *Psoralea californica* Wats.

Prostrate, near summit. ^{very}

Keel $\frac{2}{3}$ length of wings; wings about equalling banner*. Banner bluish white, wings purple. Calyx-lobes linear-acuminate (very hairy), nearly ($\frac{1}{2}$) as long as cor. Calyx-lobes 2 x as long as tube. Grows on land slips. Root very short.

* a little shorter than banner. Flowers 6 lines long.

No. 9658. *Delphinium variegatum* T. & G. Grassy open in the woods

- No. 9659. *Microseris heterocarpa*
 No. 9660. *Arctostaphylos manzanita*
 Parry. Two living shrubs and one
 dead one. Neither flowers nor
 fruits. No crown-sprouting. The
 dead shrub is dead after the
 fashion of this species. I feel sure
 from its general appearance and its
 leaves that this is *A. manzanita*;
 but neither this nor 9658 have
 been reported from south of the
 bay before.


- No. 9661. *Salvia columbariae* Bth.
 On land-slips.

- No. 9662. *Arctostaphylos auriculata*
 Eastw. This seems more like a
 cordate variety of *A. glandulosa*
 than like *A. andersonii*. It grows
 in a heavy clump, and evidently
 crown-sprouted. Obtained by Dr.
 Bode (of my party). See no. 9737.

- No. 9663. *Sanicula tuberosa* Torr.

Thin woods in loamy soil, N. slope.
 see p. 88.

- No. 9664. *Micropus californicus* R. & M.
 with 9663. I find no "scale-like
 processes on receptacle".



I find about 3
 sterile fls. in center of
 head. The basal bracts
 are scarious with an herb-
 aceous center.

- No. 9665. *Plectritis*

with 9663.

- No. 9665a. *Pedicularis densiflora* Bth. with 9663.
 No. 9666. *Thysanocarpus curvipes* var.
submarginatus ^{Japan.}
 with 9663. *elegans* Rob.

- No. 9667. *Agoseris heterophylla*
 with 9663.

- No. 9668. *Hypochaeris glabra* L.
 with 9663.

- No. 9669. *Microseris linearifolia*
 with 9663.

- No. 9670. *Sisyrinchium bellum*.
 with 9663.

- Cont. no. 9654. The achenes are strongly flattened, but flattened sideways, that is obcompressed! so that the subtending bract of the receptacle embraces the edge of the achene!

In *Wyethia*^{thin} *helenioides* the bracts of the receptacle are fully $1\frac{1}{2}$ times as long as the achenes; in *Wyethia angustifolia* the bracts are nearly twice as long as the achenes.

In *W. castanea* the bracts scarcely equal the achenes. Cont. p. 101

- Trip to Mt. Diablo with Dr. & Mrs. W.F. Bode and Miss Joyce Saunders.

The day was very beautiful. The hills southeast were in green, with fairy shades and tones coming on the shoulders. The distant hills, the whole view in fact, was drowned in a thin purple haze. We turned our heads partially upside down so as to get the advantage of the unused portions of the retina. The view was exalting!

• No. 9671. *Trifolium warrasgoldii* Lehm.
Marsh in the hills. Cont. below.

• No. 9672. *Erigeron philadelphicus* L.
Marsh, a little standing water.
Flowers white.

• No. 9673. *Agoseris hirsuta* (Hook.) Greene.
Bracts oblong-ovate, acute.
Ligules yellow, burnt umbers on back, much exceeding invol. bracts, forming a broad disk.

Grassy hills.
• No. 9674. *Agoseris plebeia* Greene.
Bracts narrowly elongate-lanceolate. Ligules yellow both sides, very short, scarcely exceeding invol. bracts.
Grassy hills.

- Cont. no. 9671. Calyx-teeth subulate, a little exceeding tube. Calyx $\frac{1}{2}$ as long as cor. Wings $\frac{7}{8}$ as long as banner; keel $\frac{3}{4}$ as long as wings. Banner whitish; keel & wings purple. Banner notched.
= *J. involucratum* Ortega.

- Cont. from no. 9655a This station is in the first cañon after one makes the turn around the point of the first ridge in going up the mt. There are some fine large sandstone rocks sticking up high out of the cañon, forming an interesting cluster. See no. 9663. See no. 9655a.

- Cont. no. 9673. Bracts dull green or dull reddish-brown, but uniform or essentially uniform in color. Peduncles in fruit not much elongated. Bracts more nearly one series than in 9674.

- Cont. no. 9674. Bracts streaked dark purple-brown and dull green. Peduncles in fruit often $1\frac{1}{2}$ times elongated. Bracts in several series.

Berkeley, June 1, 1922

- Umbellularia californica Nutt. The Botany Building is being moved. The rollers under the skids are native lumps. They stand a lot of roughage and hammering and yet are light, easily handled.

Little Arbutus Creek, Gilroy, May 29, 89

Sanita Cruz Mts.

1922

c. 600 ft.

No. 9675. *Arenaria douglasii* Fenzl.

Stamens 10, those opp. sepals the larger and bearing a scale on the back at base. Rocky outcrop.

No. 9676 *Glechoma hederacea* L.

This mint grew near Boling Ranch.
= *Nepeta hederacea* Threv.

No. 9677. *Prunella virginiana* var. *derrissa* Walp.
See p. 90. The shrubs along Moody Creek near camp showed spots of this gall. ^{cf. 9710.}

- Mr. Walker said: *Marrubium vulgare* (Horehound) and *Nepeta cataria* (Catnip): In the East these two species grow together under the same conditions. The former comes into California as a very common species in waste lands. Why does not the other one come with it? There are requirements for the growth of a plant which we do not apprehend as yet.

No. 9678. *Eriodictyon glutinosum* Bth.
 Filaments glabrous or with 2 or 3
 scattered hairs apiece. Calyx-lobes almost
 distinct. = *E. californicum* Greene.

- Dr. E. C. Van Dyke calls these gall-mites "an
 acarid of the family Eriophyidae". See his
 letter, June 7, 1922. This note refers to No. 9677.

- Miss Amy Rinehart says: The
 gooseberry in the Yosemite Region
 makes a red clear jelly that
 is very delicious. The berries are
 whipped off the bushes on to a
 sheet and cooked. When the mass
 is put in a bag to be squeezed
 out the spines go soft. [This is
Ribes roezlii Regel.]

In Napa Co. settlers mix apples
 and *Sambucus glauca* berries for
 a jelly which is superior to jelly
 made from either.

No. 9679. *Lonicera hispidula* var.
californica Climbing high
 over trees and shrubs. Fls. pink; indurated.

No. 9680. *Rubus parviflorus* var.
velutinus ^{Greene} Common in the woods.

No. 9681. *Stachys* (~~discarded~~ ^{immature})
 This mint grew along the road
 in a settled district, the spot being
 $\frac{1}{4}$ mile from a farmhouse. Summit
 of the Santa Cruz Mts. near Mt. Madonna.
 c. 1700 ft.

No. 9682. *Arctostaphylos andersonii*
 Gray. This grew along the ridge
 and formed large arborescent
 shrubs with good-sized trunks; 8 to 12
 ft. high. Near Mt. Madonna.
 1800 to 1900 ft.

No. 9683. *Fragaria californica* C. & S.
 1900 ft. Woods in opens, about
 borders of thickets, fruiting freely.
 c. 1500 ft. Near Mt. Madonna.

No. 9684. *Quaphialium*
 Near Mt. Madonna. Opens in woods.
 c. 1800 ft.

- No. 9685. *Lesqueris thyrsiflorus* Esch. Shrubs 10 to 14 ft. high are in wonderful bloom; solid masses of bloom. I also saw individuals ^{20 to 25} 22 ft. high. Near mt. Madonna. c. 1700 ft.
- No. 9686. *Oxalis oregana* Nutt. Deep shady moist woods. Gréninger Creek, tributary of Little Arthur Creek.
- No. 9687. *Potentilla glandulosa* Lindl. var. *wrangeliensis* Wolf. Gréninger Creek.
- No. 9688. *Arctostaphylos glandulosa* Eastw. Hairily crown-sprouting from heavy platforms at ground. Gréninger Creek.
- No. 9689. *Colobochortus tolmiei* H. & A. In brush Gréninger Creek.
- No. 9690. *Rhamnus californica* Esch. var. *tomentella* Gréninger Creek.
- No. 9691. *Lesqueris foliosus* Parry. Gréninger Creek.

- No. 9692. *Pickeringia montana* Nutt. white-flowered form. These shrubs are very floriferous. I have never seen these ^{series} more so. They grew on the steep bank of a cut. They arose from roots of older plants undoubtedly. The top of the bank was crowned with a thicket of the same species, the flowering blood-red! the usual color!! Mutilation of the roots - root pruning, may have produced the unusual effect in the way of abundant bloom - but why "white"? I say "white"; it was white with a flush of pink - very much like a peach blossom! Gréninger Creek.
- No. 9693. *Orthocarpus purpurascens* Bth. Gréninger Creek.

- No. 9694. *Eriophyllum emfertiflorum*

Grüniger Creek

- No. 9695. *Galium andrewsii* Gray.
 Rocky outcroppings, disintegrating rock. Grüniger Creek

- No. 9696. *Gilia densiflora* Benth.

With 9685 & 9675 Grüniger Creek.

= *Linanthus grandiflorus* Greene.

- No. 9697. ~~*Hesperolinon micranth-*~~
~~*Hemilichadus*~~ *um* (Gray) Small

This lobeloid plant grew on a rocky slope. Grüniger Creek.

- No. 9698. *Microseris douglasii* Douglasii
 Grüniger Creek.

- No. 9699. *Micropus californicus* T. & G.
 Grüniger Creek.

- No. 9700. *Convolvulus subacaulis*
 Greene. Steep sunny slope. Grüniger Creek.

- No. 9701. *Psoralea strobilina* Nutt.
 Bushy slope. Grüniger C.

- No. 9702. ~~*Gilia*~~ ^{*Linanthus*} *androsaceus* (Benth.) Greene.
 Better det'd as *Linanthus parviflorus* Greene.
 (W. L. G.). See p. 114a. Grüniger Creek.

- No. 9703. *Allium unifolium* Kell.

Fls. lavender-pink. No crests; lobes creased down the middle.

- No. 9704. *Collinsia heterophylla* Graham.
 Fls. white. Grüniger Creek.

- No. 9705. ~~*Clarkia purpurea* ssp.~~
~~*Clarkia*~~ *quadriflora*

Stach. Grassy open slope. Pts. with

- No. 9705a. ~~*Tri-lobed*~~ ^{*Tricolidatum*} *folium ciliatum* Nutt.
 with 9705

- No. 9706. *Phacelia imbricata* Greene.

= *P. calycosa* (Gray). Rocky outcrop in open.

Fls. white

- No. 9707. *Microseris douglasii* Douglasii
 with 9705.

- No. 9708. *Sidalcea malvaeflora*
 Gray. Both ♂ & ♀

- No. 9709. *Plagiobothrys canescens*

Bth. Open, trailing on ground. Corolla of achene green with a fine hollow scar in it, like an insect puncture. Achenes brown.

- No. 9710. *Prunus subcordata*
 Benth. Fruits all diseased

Little Arthur Creek, Gilroy
Santa Cruz Mts. c. 7500 ft.
Santa Clara Co.

as usual, except one. Leaves also
with another gall-type. See p. 97.
See W. T. Horne in litt. June 27, 1922.

- ✓ No. 9711. *Oenopodium californicum*

Flat

- ✓ No. 9712. *Viola ocellata* B. & G.
Shade of woods.

- ✓ No. 9712 a. *Viola pedunculata* Feg.
Little Arthur Creek.

- ✓ No. 9713. *Matricaria* ^{matricarioides (Less) Porter} ~~discoidea~~
Beaten ground near Barn.

- ✓ No. 9714. *Psilocarphus tenellus*
Nutt. var. ^{tenellus} Prostrate, on flat.

- ✓ No. 9715. *Rumex salicifolius*
Low fields near mt.

Eden.

- ✓ No. 9716.

This grass grew in a field
formerly cultivated. with 9715

(Mt. Eden in Alameda Co.) May 28, 1922.

- ✓ No. 9717. *Trifolium gracilentum* B. & G.
var. *inconspicuum* ^{Pres.} with 9715.

- ✓ No. 9718. *Avena barbata* Pott. ex Link
(Det. Beecher Crampton, Mar. 1922)

This grass grew with 9715.

— California Botanical Society.
This trip was to the camp at Balfing
Ranch, Little Arthur Creek, Gilroy.
Annual Camping Trip, Santa Cruz Mts.
I took Professor Bradley with me.
At the campfire Kellogg the Bird Man
talked, in addition to Society mem-
bers. [I talked on the plant history of the region]

- Cont. 9677. "The galls are the work of one
of the gall mites, an acarid of the family
Eriophyidae" - E. C. Van Dyke in litt,
June 2, 1922. Those on #9710 belong to
the same family.

- ✓ No. 9719. ^{Haplopappus} *(Gnaphalium) linearifolius* DC.

Making a fine show on
top of the mt. Drawing from
this. Mt. Diablo. 23 May 1922.
See p. 86.

- No. 9719a. *Convolvulus subacaulis* Greene.
Filaments narrowly linear-dilated at base, the lower $\frac{2}{3}$ adnate to the corolla-tube, each fil. just above the point of adnation bearing one slender tooth on each margin, or sometimes one margin toothless. Fils. sparingly glandular short-hairy.

I have never noticed these structures before, nor do I find them in any text; not in DeCandolle or any subsequent place. The orig. place of publ. Beechey's Voy. does not notice them. This may be a species distinct from *C. subacaulis* but it looks just the same.

North Berkeley Hills.

I have looked over plants in the field and find the fils. toothed at base. [Near the Cora

June 7, 1922

Williams Institute the plants were not toothed! This is some distance from the Blake place, where the fls. with toothed filaments are found - say a mile.) The filaments are, however, noticeably glandular short-hairy on the lower part, i.e. the adnate part.

- No. 9720. *Convolvulus luteolus* Gray.
In the open, - climbing up grasses, the stems of the year arising from prostrate woody stems, size of a short cord, and several feet long. Fils. finely glandular short-hairy.

- No. 9721. *Medicago sativa* L.

This alfalfa grew in a field near Mt. Eden, where it may have been cult., but only a few plants now present. Taken on a trip to the Butchers at Los Altos, June 10, 1922.

- No. 9722. *Eremocarpus setigerus* Bth.
Colonizing a playa on the flat
and the only occupant (living).
No. 9723. *Lythrum hyssopifolia* L.

This gives every appearance of being annual. Winter pool bird.
No. 9724. *Boissieria* ^{glabella (Nutt.) Walp.} ~~*densiflora*~~
Wats. Winter pool bird.

- No. 9725. *Ceanothus*

Vacaville, Little Oak Ranch.
This plant occurs along the
Mile Square Road as far as
the Big hill. At that point it
is spreading into the mile
square field, which in these
days is uncultivated.

- No. 9726. *Eryngium yoseyi* C. R.

Playas sw. of Vanden

- No. 9727. *Psilocarphus oregonus* Nutt.
Quite dead; on the playa
with 9722

The trees look well and show the effect
of last winter's rainfall. The buckeyes are
still rounded and green - whereas
at this time they are usually brown.
The latest tree to leaf, the first to go.

Cont. from p. 86. [no. 9654.] *H. castanea* seems
to be an excellent example of a
subspecies. I do not recall any that
better meets the case. It has not a
single technical character of separ-
ation from *H. californica*, and yet
it seems to me so distinct!
not by reason of just looking at
it, but by careful study. There
are many differences between the
two but all insignificant. It is
indeed a good subspecies.

- Mr. A. O. Garrett, of the Salt Lake
City High School, is engaged as
white Pine Blister Rust control and
assigned to Cal. for this summer. He
called this morning with S. H.

Wyckoff. He is stoutish, 5 ft. 10 in., rather dark complexioned, mustache, pleasant mannered. He said that Mormons told him that *Ribes aureum* is called Soldier Berry in Utah because the invading Gentile soldiers in early days ate the berries and became ill. Wyckoff said *Ribes cereum.* is called Squam Current and *R. bracteatum*, Skunk Currant. - June 29, 1922.

- Heermann, A. L. His three-years residence in California. Journ. Acad. Nat. Sci. Phila. ser. 2, 2: 259-272 (1853).
Cf. Grinnell, U. C. Publ. Zool. 24: 45 Also. Proc. Cal. Acad. ser. 4, 7: 335 (1913).

- Private herbaria. E. J. Durand, Prof. of Botany, Univ. of Minnesota. His herbarium just sold to Cornell Univ. (12,000 spms.). See Sci. 57: 298 (9 Mar. 1923).

✓ No. 9728. *Jaumea carnosa* Gray.
Rays 9, 9, 12.

✓ No. 9729. *Orthocarpus*

✓ No. 9730. *Limonium californicum*
Heller.

✓ No. 9731. *Cuscuta salina* Engelm.

✓ No. 9732. *Eryngium armatum*
C. + R. Adobe flats.

No. 9733. *Lathyrus vestitus* ssp.
bolanderi Summit of El Cerrito.

✓ No. 9734. *Lythrum hyssopifolia* L.
Adobe flats

No. 9735. *Centaurea melitensis*.
El Cerrito.

- Menzies, A. Cf. C. F. Newcomb,
"Plants collected by A. Menzies
on the Northwest Coast of America".
See separate. 1923.

Mt. Diablo, July 14, 1922.

No. 9736. *Arctostaphylos manzanita* Parry. This is common in the very head of Sycamore Creek cañon, behind the range of Black Hills. One can look from the the roadway across the cañon and see these bright green spots rising above the general level of the chaparral and chamise. The shrubs are 6 to 14 ft. high. The individuals are not numerous, not rare. ^{sep. 12}

No. 9737. *Arctostaphylos auriculata* Eastw. 12 ft. h. I managed to get thru the brush to a thicker of this. I found it a colony but not crown-sprouting. It was a colony of seedlings. It has a gray-green appearance at a distance. It seemed much less frequent than *A. manzanita*. Head of Sycamore Creek.

No. 9738. *Verbena prostrata* R. Cr. Cañon del Rey, Monterey.

No. 9739. ^{= Dudleya} *Cotyledon caespitosa* Haw. ^{sep. 12} Slender, tall, the tufts or rosettes of leaves small and narrow. leaves linear-attenuate, almost cord-like. Calyx lobes triangular or, better, deltoid. Petals broadly ovate, acute, distinctly joined at base. ^{sub-}filaments alt. petals with enlarged or dilated bases, far more distinctly so than in the case of the opp. fil. Corolla pale saffron. Bracts triangular-sessile. Leaves linear, acuminate, much thickened or rounded below and flat above, that is semi-biclate. Branches of the cyme distinctly one-sided. Cyme open as compared with no. 9741.

no. 9740.

*Centaurium**muhlenbergii* Wright
exaltatum Light

Sepals linear-attenuate, strongly
ridged on the back, as if conduplicate.
Corolla lobes elliptic, obtuse. Cor.
pink-red with a marked white-
eye-spot.

no. 9741.

*Dudleya farinosa**Echeverria cotyledon* N. & M. ex.*Cotyledon farinosa* Baker


Plant white-mealy, much more so
than in 9739. Calyx-lobes ovate,
acute. Calyx less syncarpalous
than in 9739. However, the dif-
ference in shape of calyx-lobes
is much more pronounced a dif-
ference. Flowers about the same size
as 9739. Corolla pale-cream-color.
Petals ovate, subacute, strongly
fleshy-ridged on middle of back,
distinctly united at base but
the degree of union very narrow.
Fleshy ridge on back no more than
in 9739. Leaves very white-floury. ^{cont. p. 107}

July 20, 1922.

no. 9742.

*Triteleia**Brodiaea ixioides*

Open pine woods, 1/8 mile from the
beach.



..... anther

Dilated fila. with
..... horns.

Cont. below.

- Cont. no 9741.

Leaves broadly oblong or oblong-
ovate, acute. Racemes of cyms very short
and infl. very compact. ^{See p. 121.}

no. 9742, cont. Per. tube just 1/3 as
long as segments. Per. segments
narrowly oblong-acuminate - or
narrow-oblong. Ovary stippled!
Per. golden with a broad band on
back of segments running down ^{brown} base
of tube, the color band narrower
below.

- No. 9743. *Carum gairdneri* H. & A. with 9743a.
 No. 9743a. *Carum Kelloggii* Gray.

Open pine stand back of Pt. Joe.

- No. 9744. *Sisyrinchium*

Open pine stand.

- No. 9745. *Elymus condensatus* Presl.
 (DET. Beecher Crampton, 1972)

This grass is 8 ft. high!

Asilomar.

- No. 9746. *Corethrogyne californica* DC.

Open pine woods back of Pt. Joe.

- No. 9747. *Astragalus menziesii* Gray.
 = *A. nuttallii* (F. & G.) Howell

Stems 2 to 2 1/2 ft. high, erect,

at length diffuse and decumbent.

- No. 9748. *Chlorogalum pomeridianum*
 (Scutell.)
 Flowers aging purple.

- No. 9749. *Navarretia squarrosa* (Esch.)
 H. & A.

- No. 9750. *Spergularia macrotheca* (Hornem.)
 Heynh.
 Pt. Joe.

- No. 9751. *Eryngium armatum* C. & R.
 Pt. Joe.

- No. 9752. *Euphorbia peplus* L.
 Monte Rey - Green's garden. He

regards it as an insufferable pest. It
 seeds freely and germinates readily
 and promptly.

- No. 9753. *Eriophyllum*

- No. 9754. *Sphaele calycina* Bth.
 = *Lepechinia*

Corolla light purple. Asilomar.
 near Pacific Grove.

- No. 9755. *Phalaris californica* H. & A.
 to 3 1/2

This grass was 2 1/2 ft. high.

- No. 9756.

This lichen grew on *Cupressus*

macrocarpa

- No. 9757. *Phacelia malvaefolia* Cham.

Under *Cupressus macrocarpa*

at the Octopus Tree.

- No. 9758. *Elymus glaucus* Buckl.

This grass grew in the open
 pine woods.

Called on Bradley Moore Davis, Prof. Bot. Mich.
 He is a tall well-formed chap physi-
 cally, good-looking, very pleasant manner.
 He said: "You are a much younger
 man than I expected to meet." Cont. p. 121

- no. 9759. *Arctostaphylos pumila* Nutt. It is distinctly mat-like and 3 or 4 in. high. The prostrate stems root at the joints or adventitiously. Occasional portions of a colony become ^{8 to} 14 in. high but the whole effect is the same - a low rough mat. Cont. p. 111.
- *Arctostaphylos vestita* Eastw. is 4 to 5 ft. high. Its bark is shreddy with the shreds persistent. This is its most marked feature. It - crown - sprouts. It looks more like *A. glandulosa* Eastw. than any other species. It looks enough like *A. glandulosa* to be a variety of it but it may be distinct. See no. 9761.
- no. 9760. *Adenostoma fasciculatum* H. & A.

[nos. 9759 to 9767 were collected east of Hotel del Monte on the sandy flats and hills]

Cont. no. 9759. *A. pumila* Nutt. On the sandiest flats, typically it forms prostrate mats 6 to 10 or 14 ft. in diameter. In better spots of soil it becomes 18 in. or even 2 ft. (!) high; but the colonies are circular low mounds and the general effect is much the same. Circular colonies joining each other will be medium-green, light green, whitish (pubescent) and various shades between.

no. 9761. *Arctostaphylos* ^{*tomentosa* Small} ~~*vestita*~~ ^{Eastw.}
See p. 110.

no. 9762. *Corethrogyne californica* DC?
This composite is abundant on the sandy flats. See p. 120.

no. 9763. ^{*Horkelia cuneata* ssp. *sericea* (Gray) Kew} *Potentilla lindleyi* Greene.
See p. 120. Not rare on the sandy flats.

no. 9764. (*Ericameria*) *Haplopappus* ~~*Eastwoodiae* Hall~~ Sandy flats. Rays 3, 2.

No. 9765. *Croton californicus* Mill.
Arg. See p. 120.

No. 9766. *Eriophyllum*

No. 9767. *Cordylanthus sylvaticus*
Jepson. = *C. rigidus* var. *argyr.* Jepson. See p. 119.

No. 9768. *Beaurothia dentata* J. & G.

No. 9769. *Hemizonia corymbosa* (DC.) T. & G.

- At Monterey I stopped with that always interesting personage, Harry Ashland Greene, who has done so much for tree-planting in California. The trees along the highway near Folsom are his Coast Live Oaks. Similarly he has supplied trees for school grounds - of course always gratis - but he found it necessary to make overtures credit- ing as to planting and care which has proved a healthy education regarding trees. Cont. p. 115

No. 9770. *Hemizonia*

gigling sta. See p. 118.

No. 9771. *Stephanomeria virgata*

See p. 118.

No. 9772. *Cirsium* ~~*occidentale*~~ ^{*coellarii*} (Nutt.)

Jepson. 2 to 3 ft. h. Frequent.

No. 9773. *Helianthemum scoparium*

Nutt.

See p. 117.

Forming a heavy tuft in a new-made sand bank.

No. 9774. *Gnaphalium*

Sandy bed of Salinas River at
Naposet.

No. 9775. *Scirpus campestris* Britt.

Low flats along the river,
flooded at high water as high
tide. Salinas River, Naposet.

No. 9776. *Scirpus americanus* Pers.

With 9775. Pure colonies over
very small areas.

No. 9776a *Leucanthemum papillosum* T. & G.
658 ft. - h.

No. 9777. *Corethrogyne filaginifolia* (H. & A.)
Nutt. var. *rigida* Gray
Scattered rather thickly over
a little hillside open, the stems
several to many in a cluster, erect.
See p. 117.

No. 9778. *Navaretia atractylodes*
H. & A. With 9777

No. 9779. *Aretostaphylos glandulosa* Eastw. *Crouseana*
tomentosa var. *cornuta* Adams

No. 9780. ^{Dubleya} *Cotyledon caespitosa*
Haw. Scattered over the low sandy
hills amongst low half-woody plants
Marina sta. See p. 117. Near Monterey.

No. 9781. (*Ericameria*) *Haplopappus*
us Eastwoodae Hall See p. 118.
Marina sta., Monterey.

No. 9782. *Artemisia pycnocephala* DC.
Sigling sta.

July 23, 1922.

No. 9783.

No. 9784.

No. 9785. *Stachys chamissonis* Bth.
Pacific Grove.

- *Linanthus androsaceus* Greene.
Cont. from p. 95, no. 9702. Mrs. Vir-
ginia Bailey determines 9702 as above.
Perhaps it were better to regard it as
intermediate between *L. androsaceus*
and *L. parviflorus*.

- Cont. from top of p. 65 nts.
These plants, nos. 9602-9614, were
collected within about 2 mi. of
Peaceful Glen, not right at
Peaceful Glen School.

- Cont. from p. 66. No. 9609. *L. nanus*
Dougl. var. *apricus* C. P. Lx. Not *L.*
micranthus

July 23, 1922.

- Cont. from p. 112 at bottom. Greene has
grown a great lot of trees of all kinds
for years. He is now quite an old man,
say over seventy, but he doesn't look
it. His father was first acting Mayor
of San Francisco (cf. Green st.). His
maternal grandfather Fish was a
manufacturer and contractor who
kept Lee's omnies going and who was
imprisoned with Jefferson Davis in
Fortress Monroe after the Civil War.
His brother is Clay Greene the drama-
tist of New York City.

Greene says Sam Tratten measured
at the ground, after falling, a Ten
Oak in the Santa Lucias, 218 ft. tall.
Also carefully estimated another
(standing) tree as 208 ft. high.

- Stations out of Monterey: [towards Santa Cruz]
Gigling sta. Salinas River Bridge
Maxima sta. (Neposet).
Bardin sta.
Lapis sta.

July 22, 1922.

- *Cupressus macrocarpa* Hartw. The thin little strip is being sold as villa lots. It is an outrage. Greene, H. F., says he counted the rings on the end of a sawed trunk as 284. He felt that he must be mistaken, since the trees are reputed so ancient, and verified the count however by going over it carefully once more.

- On my way down to Monterey I drove by San Juan Bautista; on my return I drove by the bay shore route, crossed the Salinas River near its mouth, and so on to Watsonville. Near Soquel I "chanced" into a finely wooded little hill on left side of the road. The trees were tall live oaks, delightfully spaced, with fern on the slope beneath them. Thence to Santa Cruz. Cont. p. 121.

- *Corethrogyne* (no. 9777) cont.

Several to many erect stems from the base. Rays purplish lavender, 18, 21, 21, 18. Ovaries densely hirsuticose - connescent. Sandy open hillslope. Disk yellow.

- *Helianthemum* (no. 9773) cont.

Densely branched and very floriferous. Stamens 18, 17, 16, 15. Petals yellow, roundish-rhomboidal, apiculate, $2\frac{2}{3}$ lines long. Stigma capitate; style short. Carpels 3; placentae parietal; ovules several to each placenta.

- *Cotyledon* (no. 9780) cont.

This grew on the sandy hills, covered with low scrub or nearly open in spots. The tall stems rose up grandly; they are not so very tall but they rise up stark. The plants in habit and in fls. are exactly like those at Monterey sea-cliffs. See no. 9739.

- *Ericameria* (no. 9781) ^{cont.} Rays yellow, fading nearly white or pale cream. Rays 3, 3, 3, 3, 2, 1, 3, 2, 2, 2, 1. This is the dominant species on the sandy hills for miles about Marina stn. along the coast or bay shore.

- *Stephanomeria* (no. 9771) ^{cont.}

Ligules ^{pink} purple, 8, 13, 11, 8. ^{12.} Bracts of involucre obtuse, especially the long inner ones, purplish brown, purple-dotted, ciliate at apex. Sandy hills. ^{↑ minutely}

- *Heurizonia* (no. 9770) ^{cont.}


Rays 18, 19, 21, 21, 19, 23, ^{21.} Disk corolla a little annular-enlarged at base - the annulation somewhat obscure but still an obvious enlargement, its ovary with some minute pappus bristles. See sketches with spec.

Ray achene, except base (inner), covered with boat-shaped bract.

- *Leordylanthus* (no. 9767) ^{cont.}

Inversely yoke-shaped purple in throat, below, that is on lower ^{spot} part under teeth of lower lip. Filaments inserted at summit of tube, densely hairy on one side. Stigma dilated, inversely club-shaped, bent abruptly on its style.

Sandy flats,
Monterey.



Lower calyx segment larger (i.e. broader than the upper one).

- *Patentilla* (no. 9763) cont.

Petals white, ^{narrow}obovate, drawn down to a claw-like base, $2\frac{1}{2}$ lines long. Stamens opposite calyx-segments the longer; all with downwardly dilated filaments. Stamens 10.

Petals small. "a" and "b" are mere forms of 9763.

- *Croton* (no. 9765) cont. *C. californicus* Mill. Arg. Calyx - lobes roundish-ovate, not distinct. Stamens 13, 12, 10, 11.

- ??
- *Lessingia* (no. 9762). See below

Flowers with larger 5-cleft corolla: 8, 6, 7, 5, 7. Bracts of involucre black-tipped. Fls. yellow.

Note - 9762 was a

- *Corethrogyne* (no. 9762) cont.

Rays 22, 21

Rays very pale lavender, disk yellow. Bracts of involucre glandular - pulverulent.

- *Sternis* (no. 9782).

Inner bracts of involucre with scarious margins.

 = shape of receptacle.

- On the way back from Santa Cruz I took a newly paved road that came out at the top of the ridge above Alma stn. (on the railway). It must parallel pretty closely the old road over the mts. to Los Gatos.

- Cont. from p. 109. He is teaching at the Hopkins Biol. Lab. of Stanford. He was the first man in this country to try out *Delphinium* *Amorpha*.



No. 9739

gland blind
ovary in *Echeveria*
caespitosa



No. 9741

gland blind carpal
in *Echeveria* *cotyledon*.

- *Quercus agrifolia* Neé. H.A. Greene says that there are certain trees in the Del Monte grounds that yield edible acorns - that the head gardener often fills his pockets and munches them during the day.

- ✓ No. 9786. *Anaphalis margaritacea*
 ✓ No. 9787. *Castilleja*

- ✓ No. 9788. *Ribes leptospermum* var.
 No! = *Ribes menziesii* Pursh var. *janatum* Jeps.
 ✓ No. 9789. *Actaea*

Common about the base
 of Sugar Loaf.

- ✓ No. 9790. *Pteris aquilina*
 5 ft. high.
 ✓ No. 9791. *Corylus* ^{*cornuta*, var. *californica*} ~~*rostrata*~~

- ✓ No. 9792. *Scutellaria californica* Gray
 Bushy hill slope, south exposure.

- ✓ No. 9793. *Cerasus emarginata* (Hort.)
 = *Prunus* " ^{var. *weep.*}
 Tree 18 in. circ. at 1 ft. See
 no. 9633; same tree there measured.

- ✓ No. 9794. *Anthemis cotula*

- ✓ No. 9795. *Centaurea melitensis*

- Cont. from p. 104. No. 9736. The
 rachis is darkish (or glandular) canescent,
 decidedly darkish, and rather
 tho not markedly canescent. The
 plants in the field seemed to me
A. manzanita Parry, but they
 have some points of contact
 with *A. stanfordiana* such as the
 small berries and the bright-green
 leaves. There are, however, occasional
 specimens which have trichose canescent
 and not glandular, and the rachis
 notably thickened upward, that is
 like *A. manzanita*.

- No. 9795a. *Sanicula menziesii* H. & A.
 Berkeley

No. 9796. *Anaphalis margaritacea*
♀ plants. Mostly in a clump
by themselves.

No. 9797. *Anaphalis margaritacea*
♂ plants. Mostly in a clump
by themselves.

No. 9798. *Aralia californica* Wats. cont. p. 125.

No. 9799. *Symphoricarpos racemosus*
Shrubs are now in fruit, and
some shrubs still in flower. Indeed
in one cluster I find ripe fruit
and flowers in anthesis. cont. p. 125

- *Sambucus glauca* Nutt. In full fruit
and in full flower, different branches
on one tree.

No. 9800. *Ribes sanguineum* Ph. var.
glutinoseum Wats.
Berries with rather heavy.
bloom! They are when dead ripe
soft and bland, but not at all
sweet. Cf. 9801. Foliage with a
marked aroma as nutmeg smell.

- Cont. no. 9799. Corolla distended on lower
side, that is distinctly ventricose.
- Cont. no. 9778. Leaves to 4 ft. 9 in.
long!

No. 9801. *Ribes sanguineum* Ph. var.
melanocarpum Jepson.

Berries nearly black, without
bloom! They are (when dead ripe)
soft and a little sweet. The foliage
is without aroma.

- *Cornus pubescens* var. *californica*
Berries in the shade turning
bluish; otherwise ^{clear} ivory color in the
sun.

No. 9802. *Ribes leptospermum* Cov.

No. 1 = *R. menziesii* Pursh var. *faustum* Cov.

No. 9802 a. *Urtica holosericea* Nutt.

Aug. 23.

No. 9802 b. *Senecio hydrophilus* Nutt.

Aug. Arlington, N. Berkeley Hills

Discarded
Badly damaged by
insects. L.H., 1965

No. 9803. *Minulus moschatius* var. ^{longiflorus Gray, det. A. S. G.}
 Corolla 10 to 12 lines long, 7 or 6 lines broad. Lobes of the lips much alike, short and broad, transversely elliptic-oblong, very obtuse or obscurely apiculate*. Stigmas thin, (sometimes) rose-denticulate. Folds of throat densely covered with very straight and stiffish hairs. Branching, but the branching long and weak. Cor. bright lemon-yellow. Ovary ^{oblong} a little obcompressed. - Moist-springy place, Inverness.

No. 9804. *Galium californicum* H. & A.
 Trailing over the floor beneath the alder and buckeye woods. Cor. lobes with a very short but very marked bristle tip. Fls. white, very small but when scaled ^{dull} 1 1/2 lines broad. c. 20 ft.

* sometimes a little notched.
 Cont. p. 128.

No. 9805. *Stachys chamissonis* Bth.
 This becomes 6 to 8 ft. high! These stems are unbranched! at least short of the inflorescence. Sometimes, indeed, the stem is wholly unbranched, sometimes it bears three about equal "racemes", a terminal one and two lateral. The odor is very strong and very disagreeable; it is only with some difficulty that one can establish the basic mint odor. The odor is not only disagreeable but very clinging; it sticks on my clothing for hours after we leave the spot at Inverness where we found it - so abundant. c. 20 ft.

Flowers 11 to 16 lines long. Galea obscurely retuse, pubescent on back with ^(quite entire or) gland-tipped straight hairs. Corolla-tube

with a zone of short but densely set hairs ^{to 2} $\frac{1}{2}$ lines above the base, interrupted on the upper side opposite the style but not completely interrupted, that is the hairs very much thinner and fewer at that point.

Middle lobe of upper lip rounded, cupped, ascending, lateral lobes a little turned downward. Cor. tube below hairy band white or whitish. Cor. about twice as long as calyx.

Style glabrous.

Flowers 4 to 6 in a whorl.

Upper pair of filaments flattened, hairy on the edges; lower pair of filaments flattened, hairy on the upper edge.

-Cont. from no. 9803. The corolla, tho the lobes are subequal, is distinctly 2-lipped, due to the position of the lobes and to the prominent hairy palate.

No. 9806. *Helenium puberulum*

Achenes hispidulose on its nerves;
 ↑ dark brown,
 paleae 5, tipped with an awn or subulate point a little longer. Achenes a tuft over $\frac{1}{2}$ line long. Rays 3, 4 and 5-lobed. 3-lobed rays with 2 style-branches; 4 and 5-lobed rays with 3 style branches.

Laquearia

No. 9807. *Baccharis douglasii* DC.

Stream banks near Fairfax.

I notice only ♂ plants.

No. 9807a. *Rhamnus californica* Engelm.
 Near Fairfax. ^{typical CALIF.}

No. 9808. *Convolvulus luteolus* Gray.
 Climbing over bushes on stream bank. Near Fairfax.

- Miss ^{Ellice} Eastwood came to Berkeley a few days ago but I did not ~~have~~ to see her unfortunately. She still holds on at the Academy of Sciences in San Francisco in spite of her enemies. It is about two years since Hall told me of the "problem" of Miss Eastwood. He is now on the council of the Academy and responsible for the botany interests as discussed in the Council. He says, Miss Eastwood goes out and collects wholesale, getting as many things as she can, breaking off a twig here and grabbing a plant there - but making no data or real field studies to go with the plants. Building up a large herbarium as rapidly as possible but paying no attention to quality.

Similarly after the fire an attempt was made by the

Academy people to get rid of her but she stuck. As she told me about it she jumped upon a bench, folded her arms defiantly, threw back her head and with flashing eyes proclaimed: "They'll have to drag me out by force and I help them so!" Miss Eastwood is a fighter. Her militant personality is however only known to her enemies for the most part. To most people her bright vivaciousness and cordial humor make her a well-liked person.

No. 9809. *Convolvulus arvensis* L.
Berkeley, street parkings.

- Haenke, T. He edited the eighth edition of Linnaeus, *Genera Plantarum*. Vindobonae, 1791. I note this from Britzel, *Thesaurus*, ed. 2, p. 189.

- went to Bay Farm Isl. with grad^{uate} students, namely Dr. Houghton, Miss Thacker and my artist, Miss Saunders.
 No. 9809a. *Frankenia grandifolia* C. + S.
 No. 9810. *Suaeda californica* Wats.

We find a number of plants at the old-time station, just beyond the bridge from Alameda on the lower edge of Bay Farm Isl. The plants are semi-prostrate, that is forming mats 6 ft. by 6 ft. about with the ends of the branches ascending.

- No. 9810a. *Striplax*.
 No. 9810b. *Salicornia ambigua* ^{*S. virginica* L.} Michx. ♂ & ♀
 See p. 135.
 No. 9891. *Tetragonia expansa* Mun.

Stamens 16, 12.

Calyx with one broad lobe on the upper side and 4 unequal lobes on the lower side of the flower. Stigmas 6, 7. This occurred

Sept. 9, 1922

in very few individuals along the levee.

— Mr. Charles Nordhoff, is here in Berkeley, stopping at the Berkeley Inn. He says the *Jajoba* bush* is common and a great resource to the natives in Lower California. The nuts are eaten roasted. Roasted they make a drink similar to chocolate. The bush grows in the most arid districts. It is said of it that it requires only one rain^{in a season} to make it bear fruit. It is browsed heavily by browsing animals. When such is the case it does not bear. It is cultivated in the Argentine where there are superior varieties it is said. — Oct. 17, 1922.

* *Simmondsia californica*

- No. 9811a. *Lavatera assurgentiflora* Kell.
 Prob. more escape. Bay Farm Isl.

— *Lactuca* is more
or less common thru the cañon,
both on open slopes and in the
chaparral. See no. 9819.

No. 9812. *Helenium puberulum*

No. 9813. *Habenaria elegans*

A single individual.

No. 9814. *Aster radulinus* Gray

No. 9815. *Rosa gymnocarpa*
Nutt.

No. 9816. *Quercus morehus* Kell.

A fine compact small and
sharply defined thicket of it—
8 ft. high on Pigmy Ridge.

No. 9817. *Quercus wislizenii* A.
Dc. with 9816. var. *brutescens* Engelm.

No. 9818. *Salix* ^{*serotiana* Bunge} ~~*flavescens* Nutt.~~

Near Sugar Loaf.

No. 9819. *Lactuca serriola*

No. 9819 a. *Dica occidentalis* Gray.
Upper slopes beyond Sugar Loaf.

— Elmer → Applegate, (Klamath Falls,
called today. He said: "I have never
seen *Cercis occidentalis* in southern
Oregon nor *Pinus sabiniana*. Swarth
of Univ. of Ore. refers to *Pinus*
contorta in sw. Ore. Fiedler's Gulch
[of your Silva] is a branch of Jose-
phine Creek. I have myself seen
Picea breweriana at the head of it,
just west of Kerby." Cf. vol. 20: p. 73
of this F. B.

[*Salicornia ambigua* Michx. Cort. from
9810b. The plants are dioecious, as I
find them here. I see none but
pistillate fls. in pistillate spikes and
none but staminate flowers in stamin-
ate spikes.

No. 9820. *Bonium maculatum* L.

No. 9821. *Baccharis douglasii* DC.

The colonies of male and female plants keep rather separate. The female heads tend to be rather bluntly pointed and are marked by the styles, which soon turn black. The heads of ♂ fls. are rather bracteate, since the expanded corollas make them so. The corolla in *B. douglasii* is remarkably short, filiform and without obvious limb but "toothed" very decidedly at the limb or margin, or perhaps roughly hispid rather than toothed. Low moist spots.

No. 9822. *Picris* → *echinoides*
glabrous, erect.

No. 9823. ~~*Solidago*~~ *occidentalis*
(Nutt.) T. & G. Low moist lands.

No. 9824. *Heliotropium curassavicum*
var. *oculatum* (Heller) Ten.

L. The ovary is slightly 4-creased longitudinally and thus becoming 4-lobed.

— *Prunus ilicifolia* Walp. A tree at H. A. Dutton's place, Los Allos, is 30 ft. high. It presents a fine sight now, the crown covered with racemes of large shining red cherries. The fruit is abundant, looks very enticing, tastes most excellently, but alas! is nearly all stone! The seeds germinate readily in broken ground, that is when self-sown and seedlings are abundant says Dutton. This is not a planted tree.

No. 9825. *Prunus ilicifolia* Walp.
Los Allos. Fruits. See above.
Santa Clara Co.

Mtall, Fls. of Pitt. 1:244. Century Mag.
32:238

Sept. 27, 1922

- *Lychnis rostrata* One bush
15 ft. high!; another = 19 ft. h!
No. 9826. *Chrysopsis villosa* var. *bolanderi* Gray. Rays 11, 10, 10.
No. 9827. *Zauschneria californica* Presl.
Grassy slopes.

No. 9828. *Solidago*

Margin of brushy slope.

No. 9829. *Scutellaria californica* Gray

- Coyotes. Forest Rangers killed 274 mountain lions, coyotes and bobcats on National Forests in Cal. in 1922. Santa Barbara Nat. For. land with 105 coyotes, 53 wildcats and 41 mountain lions. - District Five News Notes - about Jan. 1923.

No. 9829a. *Eriogonum caninum* (Greene) Munz
Oakland Hills, Sept.

- Menzies, A. was also an artist; see his botanical drawings in Hook. Bot. Misc. 1: pl. 23, 24, and in Hook. Fl. Bor. Am. 2: pl. 148.

- No. 9830. *Mentzelia laevicaulis* T. & G.
Biennial. Stream bed White Sulphur Creek.
No. 9831. *Brassica adpressa* Boiss. ^{= *B. geniculata* [cont. p. 140.]} 2 ft. h.
~~incand. Meigen~~ Biennial certainly; apparently perennial!! White Sulphur Creek bed. 2 ft. h.
No. 9832. *Tamarix gallica* L.
White Sulphur Creek bed.
No. 9833. *Chrysopsis*
2 1/2 ft. h. bed of White Sulphur Creek
No. 9834. *Zauschneria californica* Presl.
Bed of White Sulphur Creek
No. 9835. *Chenopodium* ^{*ambrosioides* var.} *anthelminticum*
Bed of Napa River.
No. 9836. *Quercus lobata* Nutt.
Sep. 14. Mouth of Corn Creek. See p. 140.
No. 9837. *Helianthus scaberrimus*
Stream bed, near mouth of Corn Creek.
No. 9838. *Nicotiana bigelovii*
Wate. Stream bed near mouth of Corn Creek.
No. 9839. *Eremocarpus setigerus* Bth.

No. 9840. *Salix laevigata* Bebb.

This species forms a dense regular line or screen along Coun Creek in the valley opposite Rutherford. The row is quite splendid, and I have indicated its exact location on my Napa Quadrangle. Trees 60 to 75 ft. h.

- *Sequoia gigantea* Dec. An individual stands at turn of road, right angle turn east, just before reaching willow line no. 9840.

- "Mouth of Coun Creek"; by this is meant point where Creek leaves gorge of Sage Cañon and runs over valley floor. Better say "mouth of Sage Cañon." Coun Creek mouth is down the valley near Yountville.

- *Mentzelia laeviculis*, ent. 9830. The stamens are in several rows and inserted on a narrow disk or border to the summit of the ovary. This disk appears ^{flat} like "calyx" tissue, or receptacle tissue than like an extension of the

flat ovary summit. The petals are inserted immediately behind them and quite free from them.

- cont. from p. 139. no. 9836. Acorns remarkably thick! See box material.

No. 9841. *Vitis californica* Bth.

St. Helena at the ^{river} stone bridge

- from Nelson, Professor of Botany at Wyoming is here on a vacation. He arrived about Friday, Nov. 17, stayed for about twelve days. I invited him to speak before the Seminar and he consented to say a few words, Nov. 27.

- Carl Skottsberg, Director of the Botanic Garden at Gothenburg, Sweden, lectured before the Cal. Bot. Soc.

Nov. 16. Setchell gave him a dinner that evening and invited Miss ^{Rice} Eastwood, Mr. & Mrs. ^{S. B.} Parish, Dr. & Mrs. B. W. Evermann, Dr. & Mrs. ^{A. D.} Houghton, Mr. Waltzinger and about fifteen more. Skottsberg (cont p. 142)

Shigenori Kawagoe:
The Kagoshima Imperial
College of Agriculture and
Forestry, Kagoshima, Japan.

This Japanese professor came to call this morning and brought a lot of Californian plants to name. A considerable number of them are scraps and fragments he had obtained from Mrs. Rice's Wild Flowers show last spring, and were ~~written~~ to name. He has been collecting in addition on his own account and had my Flora of Western Middle California which he seemed to value greatly. He also brought a copy of my Silva in order to

have my inscription in front.

cont. from p. 141.
said of O. Warburg, the Begonia specialist: "He is doing much better now that he is away from Berlin and Dr. Engler". This is interesting as a light ^{TA.} on the attempts to make all scientists organize under a directing head. I myself was not at the dinner*. Professor and Mrs. ^{Richard M.} Holman, Professor and Mrs. ^{N.T.} Gardner were undoubtedly there.
* See p. 203.

- Walked to Glory Hole rim, Oakland Hills, Dec. 17. Along the Sacramento Short Line Tracks *Brassica adpressa* is starting afresh from the plants of last summer, leafing out from the old stems at 1 and 2 feet. So also new shoots from *Picris* *Echivoides*. Along the wet shore

of Lake Temescal I collect:—
No. 9842. *Artemisia biennis* Willd.

It is in dense growth
right along the water margin.

— Brandegee, Mrs. K. From conversations with her, Mrs. S. B. Parish infers that her first husband, one Curran, was a rough Irish miner in the foothills of the Sierra, in one of the mining counties (probably Eldorado County) and that he was ill a long time before he died. (The "dying" is not clear.) He may have been divorced from Mrs. B. it is conjectured. At any rate freed of Curran she came to B.F. and took the course and degree in the med. dept. of Univ. of Cal. Little is known about her early life. I remarked to the Parishes, that for a woman as unrestrained and reckless in her comments on other people as

Mrs. B, she was singularly reticent about herself and particularly about her early history. — Dec. 24, 1922.

— Cleveland, Daniel. Mrs. Parish says of him: He gave up botany long ago. He had sickness, broke his leg and so on. However, he maintained sufficient interest in botany to borrow money from the Brandegees to meet his needs. Mrs. B. said he never repaid it but she took the better part, the more valuable portion, of his herbarium in payment and also some books.

(partial) Cleveland has rather recently married. — Dec. 24, 1922. See p. 201.

— San Diego Society of Natural History is, says S. B. Parish, one of the oldest organizations on the coast. Some one gave it a well-situated lot on which a hotel was built, the hotel paying ground rent and later falling to the Society. The

society required one of the Panama-California exposition ^{Buildings} at the Balboa Park, San Diego, for a museum.

Parish examined the very small herbarium in 1919 and saw nothing of value therein. - Dec. 24, 1922.

- Orcutt, B.K. Drove the team, (for hire, being employed for that purpose), which took Parry and M.E. Jones on a trip to Escondido from San Diego. On the way back they camped at Tia Juana. The next morning Jones proposed to continue the journey back to San Diego. Orcutt explained that, it being Sunday, his religion prevented his traveling on that day. Jones pulled out a pistol and forced Orcutt to go ahead. Parish thinks Pringle and Parry both were members of the party and that they had somehow

separated from the others before this incident. Parry was certainly on the expedition, but I never understood before that Pringle was of it. At any rate this was Orcutt's first botanical experience and he took so much interest in the activities of the party that Parry encouraged him to develop an interest in botany. Orcutt is certainly the promoter of ^{unique as} shady botanical schemes, the promoter and organizer of semi-swindling devices. He is a sort of botanical confidence man. His latest plan is a hotel for retired botanists where by taking so much stock you can live, ^{i.e. reside,} for so long. This ^{verbally} noted from S.B. Parish. - Dec. 24, 1922.

- Went up Tonalpays Jan. 7. from Ross Valley, coming back to Corte Madera.
- *Arctostaphylos montana* Eastw. Spreads out over the ground, forming rough mats 10 ft. x 5 1/2 ft., along

148 Mt. Tamalpais, Jan. 7, 1923.

The Eldredge trail wherever there is an "open" in sandstone rock. The ascending branches are not over 4 to 6 in high. Much like, in this respect, the more prostrate forms of *A. pumila* Nutt. at Monterey.

✓ No. 9842. *Artemisiabiennis* Willd.

Very common in all the lower woods, the most common fern.

✓ No. 9843.

A frequent fern in the lower woods.

✓ No. 9844. *Ribes malvaceum* Sm.

Las Trampas Ridge, Feb. 18.

Contra Costa Co.

✓ No. 9845. *Pedicularis densiflora*
B.R. Parasitic on *Aclenostoma fasciculatum* root $\frac{1}{2}$ in. in diameter! the union complete and striking.
with 9844.

Berkeley, Jan. 8, 1923. 149

-(Marihuana?) This is a drug obtained from some Mexican species of *Stragalus* ^(?) says Dr. A. S. Houghton. It is imported and used by Mexicans at Los Angeles. They smoke the leaves or herbage and throw them into a state of great ^{the drug} excitation, physical excitation - it being a motor stimulant. Under this stimulant they may, for example, run head first at a wall. Their physical excitement is so great as to cause cutting right thru the flesh to the bones of the manacles put on hands or feet. One of the curious phases is that after coming out of this state, the addict declares he has had a wonderful time, a splendid time! Dr. H. has promised to secure botanical specimens. I believe the Mexicans use native species when they cannot get the imported herb.

- Drove to Stanford, took dinner with the Percies, Feb. 4, went on down to Los Altos to see Sutton. He said: "When I was a lad, in the early eighties, I went to Los Angeles. It consisted mostly of adobe houses with some frame houses. *Brassica rugia* covered the whole country about. It was 10 to 12 ft. high, with stems nearly as thick as your wrist, and we children did not dare wander into such a forest because we should have been lost, as one could not see out."

- Under side of ^{bark} scales in *Pinus ponderosa* = yellow bloom; and in *Pinus jeffreyi* = chocolate. That is scales of bark! J. O. Martin, entomologist, says these are differences marked at ^{V.S.} by Brown, my student at Eagle Lake. *Dendroctonus jeffreyi* is a beetle

found only in *Pinus jeffreyi*; whereas *D. brevicornis* works on *P. ponderosa*. The latter insect is more catholic in taste; but *D. jeffreyi* is found only on *Pinus jeffreyi*. This acc. to J. O. Martin who came in to call Feb. 13.

- *Sequoia gigantea* has resin ducts as follows: peduncle, scales, axis of (woody) cone; in first annual ring of wood in vigorous branches of adult trees, none in young trees; in leaf trace of very vigorous leaves of adult trees.

Sequoia sempervirens; none in cone; none in branches or leaves.

In both species appear as a result of wounds.

- From Jeffrey, Bot. Soc. Nat. Hist. Mem. 5: 441-459 (1903).

- C. G. Pringle, & other collectors and cultivars. See Bailey, Cyclop. Hort. ed. 2 under Horticulturists.

Mt. Diablo, Mar. 11, 1923.

- ✓ No. 9846. ^{= *Microseris*} *Julia gracilis* Hook.
South slope, amongst oaks,
Black Hills. c. 1300 ft.
- ✓ No. 9847. *Ceanothus cuneatus* Nutt.
Black Hills. c. 1300 ft.
- ✓ No. 9848. *Dodecatheon Hendersonii* Gray.
with 9846. c. 1300 ft.
- ✓ No. 9849. *Viola pedunculata* T. & G.
Half-way up south slope. c. 2250 ft.
- ✓ No. 9850. *Cardamine* ^{= *Dentaria californica* Nutt.} *integrifolia* var. *californica* Japon.
Oak-covered open slope, more or
less in shade. Half-way up. c. 2250 ft.
- ✓ No. 9851. *Pinus sabiniana* Dougl.
Black Hills. cone. c. 1300 ft.

- H. E. McMinn, Professor of Botany
at Mills College says: One of my
students who was in Abrams
laboratory at Stanford tells me
that Abrams said: I am disgusted
with my account of *Arctostaphylos*
in the North American

Berkeley, Mar. 31.

Flora". This remark was apropos of
my treatment of the genus in my
recent ^{student} *Madroño* where the *Knuth* the pins
out from under Abrams treatment
completely, tho of course I do
not say so, nor even intimate it.

✓ No. 9852. *Scaevola spicata* var. *arguta*
Torr. Strawberry Cañon. Apr. 12.

(Discarded, insect damaged)
✓ No. 9853.

Along Oakland - Antioch track.
Apr. 14.

✓ No. 9854. *Orientalis europaea* L.
~~var. *retifolia* Hook.~~
Oakland Hills. Apr. 14.

✓ No. 9855. *Laetia glabrata*
Alviso, Apr. 15.

- Went today to the dedication of
the Jepson Tower (by Cal. Botanical
Society) at Lake San Andreas. The
exercises were simple, dignified and
impressive. W. W. Macmillan, Master of
the Day, Frank L. Dodd and W. F. Bode
and Mrs. A. Frederick, speakers.

= *Tritelia*✓ No. 9856. *Brodialoa laxa*

Adobe hillsides, among wild oats.

There was a great abundance of this on the lowest slope of the foothills, south side of the mt. It was scarcely as tall as the wild oats, but the effect as the wind swayed the oats was wonderful in the extreme. (Cont. p. 158)

✓ No. 9857. *Trifolium fucatum* Lindl.✓ No. 9858. *Trifolium gracilentum* T. & G.✓ No. 9859. *Orthocarpus purpuscens* Bth.✓ No. 9860. *Trifolium albobipurense* T. & G.✓ No. 9861. *Achyrochaena mollis* Sch.✓ No. 9862. *Geranium dissectum* L.

Shade of oaks. Claws markedly ciliate above the base. Styles 5. Stamens 7 to 10, filaments dilated. Cont. p. 158.

✓ No. 9863. *Torilis nodosa* Gaertn.

Shade of oaks

✓ No. 9864. *Evax sparsiflora*

Steep adobe bank in foot-hill cañon. Sterile fls. 3, 4.

✓ No. 9865. *Anagallis arvensis* L.var. *coarctata* Ledeb.

Corollas royal purple, no red ones in the colony. Colony on open hillside in little foothill cañon, about 12 ft. square. A large number of plants. (Cont. p. 158)

✓ No. 9866. *Lythia helenioides*

Rays 15.

Not abundant. Higher slopes, middle alts. on the mt. and below

✓ No. 9867. *Pterostegia drymarioides*

No. 9867a. Shade of oaks.

✓ No. 9868. *Nemophila*

Corolla clear white, large, rather showy, making masses in deep shaded cañons.

No. 9869. *Dryopteris arguta* (Kauff.) Watt
(Det. Tryon, 1958)

This fern grew on deep
cañon side, wooded.

No. 9870. *Sanicula bipinnata*
H. & A.
with 9869

No. 9871. *Ranunculus californicus*
Benth.
with 9870

No. 9872. *Plantago*
Open hilltops.

No. 9873. *Othocarpus attenuatus*
Gray. with 9872.

No. 9874. *Baeria*
with 9872

Rays 11, 11, 8.

No. 9875. *Othocarpus pusillus*
Benth. with 9872.

No. 9876. *Pinus attenuata* ^{*attenuata* *lambertii*} ~~*fusculata*~~ ^{*lambertii*} ~~*montana*~~

This tree grows on the
rocky ridge described on
p. 159. It was the nearest
and most isolated tree
but was difficult to

reach on account of chaparral
even when within 10 ft. of it;
the problem was baffling. I
had neglected to bring my
belt axe — which makes
such a difference in chap-
arral country.

— This high country is very delightful. It
is a "brimming sun-bathed world" for
one thing today; and the greens are wonderful.
^(spring)

No. 9877. *Micropus californicus* F. & M.
with 9872

No. 9878. *Micropus californicus* F. & M.
with 9872

No. 9878a. *Plagiobothrys* ^{*notofulvus* Gray} ~~*caescaus* Bth.~~

No. 9879. *Phacelia distans* Bth.

Foothill cañon, wooded.

Fls. white or nearly white.

— *Heracleum lanatum* Michx. In deep
gorge Syomure Cañon; cf. p. 159.

No. 9880. *Microseris douglasii* Douglasii
Foothill cañon, not
wooded except sparsely.

- ✓ No. 9881. *Microseris douglasii douglasii*
with 9880.
- ✓ No. 9882. *Microseris douglasii douglasii*
with 9880.
- [No. 9865. Cont. Petals crenulate.
cont.
[No. 9862, Filaments ciliate at the
middle. Petals with a white tuft of hairs
on each side above claw.
- ✓ No. 9883. *Lasthenia glabrata*
Rays 12, 14, 12, 12, 13.

- Cont. from p. 154, 9856 - I went
out to the mt. by electric train
to the station at Diablo, walked
out the Black Hawk Road, then
cut across the fields over the
low rolling hills into a cañon
leading up to "inspiration
point" on the mt. road which
here encircles the Black Hills
and then turns westerly on its

way up the mt. I plunged down
then the gap below the point into
Sycamore Cañon, - a lovely deep
wooded ^{and wild} gorge, - followed up its
east fork, on, on, and up, up -
until reaching the high lovely
opens and swales bordered by fine
oaks and buckeyes at its very
head, ^{my objective (and central station.)} Here one is near and can
easily survey the high and
picturesque rocky ridge which
circles this station on the north
and east. The ridge is very
broken and rocky, covered with
chaparral, and has two fine
clusters of *Pinus tuberculata*
on high points across the
cañon eastward. The trees
seemed as much as 40 ft. h.,
but may have been higher.
A few trees are scattered along
the ridge westward. See #9876.

Berkeley, Apr. 28, 1923.

- Redwood, no. of seed for pound,
124,900 to 224,000.
- Seed of germination, 20 to 36.
- H. C. Tott, master's thesis.
from W. L. J.
- Lemmon, J. G. See letter ^{from W. L. J.} to C. H.
Shinn, Apr. 28, 1923. A. Kellogg also
mentioned.
- Climate, - Rains early and heavy
all winter up to end of Feb. The
precip. in Feb. light. First 30 days of
March no rainfall at all! Slight
fall on last day of month, .03 inch.
March, 1923, therefore a record. An
excess of north winds in ^{new} Mar. April
rainfall heavy for April. - {Apr. 29.}
- Menzies, A. The barren-ness of his
journal in botany. See letter to
C. F. Newcombe, 8 May 1923, from
W. L. J. - And yet for introduced things in
Europe it is Menzies & Douglas & others.
- Botanical exploration. of, Chas. W. S. Smith,
Checklist of Books and Pamphlets
relating to the Pacific Northwest.

St. Helena, Apr. 29, 1923 161

- ✓ No. 9884. *Erodium botrys* Bertol.
Nearly erect, and taller than
I have ever before seen it.
Hill slope, below channisal, in
open ground.
- ✓ No. 9885. *Oenothera ovata* Nutt.
Open ground, valley swale at
edge of foothill.
- ✓ No. 9886. *Filago gallica* L.
With 9884.
- ✓ No. 9887. *Filago californica* Nutt.
With 9884.
- ✓ No. 9888. *Micropus californicus* F. & M.
With 9884.
- ✓ No. 9889. *Eriophyllum*
Borders of channisal, on
steep slope.
- ✓ No. 9890. *Girsium*
Crimsa fl. with 9889.
Branching from base, 3 1/2 ft. h.

✓ No. 9891. *Microseris linearifolia*
with 9889.

✓ No. 9892. *Wyethia*

In a little flat on border of
the foothills. Rays 13, 13, 13,
14, 16.

✓ No. 9893. *Silene gallica* L.
with 9892.

✓ No. 9894. *Platystemon californicus* Bth. with 9892.

✓ No. 9895. *Thysanocarpus curvisper*
Hook. var. with 9892.
↑ *Telegam* Rob.

Nos. 9884-9895 were collected
on a hillside just above the
valley level a little north of
the ^{valley} mouth* of Coun Creek, or
on the flat below it. See also

No. 9899. *i.e., mouth of Sage Cañon.

— My brother-in-law, Frank Pellet,
says he can distinguish between
songs of every individual lark
of the various ones that sing in the
fields and on the fences and tree
tops near his home - especially in
the early morning when he lies
abed listening to them.

Vanden sta. Apr. 30.

✓ No. 9896. *Lilaea subulata*
In pools e. of Suisun.

The ♀ spikes appear to contort and
get down under the water after
anthesis.

— The flats south of Vanden are
now having their short hey-day of
beauty. The playas are a pale sea-
green with a dense short growth of
Elygium roseyi, bordered by a ^{young} zone of
Baccharis 9898; an embroidery of deep
gold. Likewise there are gold med-
allions of the same scattered over-

30 Apr.

No. 9897. *Psilocarphus ~~argutus~~ ^{mutt.} brevissimus*. In bed vernal pool,
near Vanden sta.

No. 9898. *Baeria*.

Borders of playas a little
west of Vanden sta.
↑ of south of

over the grassy fields, relieved
by the shining or silvery heads
of *Achyrachena mollis* - now
just bursting into its fruit-
ing heads.

The wheat fields look
well, exceptionally well -
but the outlook was desper-
ate one month ago today
after 45 days of drought.

No. 9899. *Plantago bigelovii* Gray
terete. Vanden sta. Leaves semi-

- Mrs. ^{Louisiana} Scott, who is working
with a hygienic company at
West Berkeley, says: Dr. Harry E.
Alderson has perfected an
extract of *Rhus diversiloba*, alcoh-
olic, which is administered suc-
cessfully in hypodermic injections.
↑ graduated

Dr. Alderson's office is 240 Stockton st.
This product is put on the market
by Broemmels Pharmacy, 2501
California st., S.F.

- Mule-Tail or Mare's-Tail. Possibly
Equisetum sp? Pest in garden at Tib-
uron. (Loc. R. Press, 7/14/'23. "W.F.T.")

No. 9899a. *Brodiaea*

With no. 9895. This has a
couple of points or projections
to filament above insertion on
anther, on back of anther. See no.
9981.

Berkeley, May 8, 1923.

No. 9900. *Myrica californica*
 Strawberry Cañon. ♀ shrub
 with some ♂ catkins.

No. 9901. *Myrica californica*
 with 9900. ♂ shrub, with
 ♂ fls. only acc. Herbert L.
 Mason, Bot. 112.

(Discarded)
 No. 9902. *Heuchera micrantha* Dougl.
 Strawberry Cañon; foliage
 from one plant.

(Discarded)
 No. 9903. *Heuchera micrantha*
 Dougl. Foliage from one plant.
 With 9902.

- Martindale, Lustin E., 1127 Castalia
 av., Los Angeles, comes in with
 some very remarkable flower
 photographs colored. Rather finer
 than anything I have seen
 before.

Berkeley to San Andreas

25 May 1923.

No. 9904. *Lemna*

Middle River, San Joaquin Co.

No. 9905. *Gratiola ebracteata* Oth.
 c. 650 ft. in muddy bed of former vine
 pool. Valley Springs, Calaveras Co.

No. 9906. *Gnaphalium*

with 9905, on dry bank.

No. 9907. *Filago gallica*

with 9906.

No. 9908. *Brodiaea* ^{*elegans* Hoover}
~~*coronaria* Jepson~~
 det. Hoover.

Stylus approx. in center around
 style; staminodia exert against
 perianth. With 9906.

J. A.
 - Cont from p. 170. Judge Smith de-
 clares there was no Four Coun-
 ties Courthouse! See his letter
 dated June 26, 1923. I wrote him
 in order to verify the matter.
 [In Bancroft's Handbook Almanac of the
 Pacific States, p. 103 (1864), Eldorado, Ama-
 dor and Calaveras appear in one judicial dis-
 trict.]

No. 9909. = *Navarretia* ^{*gila*} *intertexta* Stand.
 Cor. very pale lavender (almost white); stamens equally inserted in tube. Fls. $4\frac{1}{2}$ lines long. Calyx-teeth very unequal, 2 long and 3 short. Leaves pinnately parted with filiform divisions, the divisions entire or sparingly toothed. Calyx-teeth with a tuft of hair on the base of the lobe, inside, otherwise glabrous or nearly so; calyx-tube pubescent outside, except at very base. All the fls. examined showed 2-cleft styles. With 9906. ^{shortly}

= *Navarretia intertexta* Hook. (No. 9909).

-Teratology. - No. 9909. One fl. with 6-merous circles, but only 2 branches to the style! It is remarkable that increase in number affected all circles except the gynoecium!

- A Blue jay sails along the hill slope above Colavertis Creek. We sit on the limestone outcrops above and watch his supremely graceful flight with saucy bows and swells in his very flight itself. The reason man doesn't like the Blue Jay is exactly the same reason that he dislikes the Coyote. These animals are too smart for him. The Coyote beats him at his own game. The trait of man is cunning. The Coyote beats him. The trait of man is deceit. The Coyote beats him. The Blue Jay is saucy and impudent. Man doesn't like the Blue Jay. He wants a monopoly of those traits. Now a small boy likes a Blue Jay. A small boy is unprejudiced; he is frank and honest and normally tells the truth - especially where others are concerned.

- The Four Counties Courthouse stood in Old Gulch, on a little ridge, maybe twenty feet above the flat. It had entrances at opposite ends. At the west end stood a fig tree; at the east end an oak. If the jury left the building by the fig tree end it was understood the accused was held innocent; if by the oak end the verdict was guilty. Justice was certain and prompt in those days and the jury acted as executioners and strung the guilty man up to the oak tree. The fig tree still stands; it is plainly an old tree tho' not a large one. The oak tree still stands; it is not a large one (being 34 feet high) but shapely, with a trunk 2 feet in diameter at 4 ft. It is an Interior Live Oak (*Quercus wislizenii* A. D. C.). My measurements are by the eye only. — Cont. p. 167. See also p. 201.

- Old Gulch lies on Calaveritas Creek, some six or seven miles S. from San Andreas.

[Cont. no. 9954, *Brodiaea*, from p. 191. The gravelly soil is so hard, so sun-baked, and more especially the corns are so deeply seated (8 to 9 in.) that it was a truly difficult task to lift the corns uningured or get them at all. The feeling was that I had set myself in for an all day's steady "pick and shovel" kind of job with just a small botanical pick. The below-ground stem and corn are very heavily sheathed in old coats and fibre. I know no other lily so heavily sheathed except (cont. p. 174).

[Cont. 9960. This is as near H. elegans Jepsen as could be expected, I think, save only in color of flowers. I had misgivings as to this color diff. which came with Tracy's spurs and notes and now I feel that the diagnosis should read in Fl. Cal. 330, fls. whitish or greenish.

No. 9910. *Calochortus venustus* Dougl.
Comet Mine, San Andreas

No. 9911. *Orthocarpus densiflorus* Bth.
With 9910.

No. 9912. *Myethia angustifolia*
With 9910 (DC.) Nutt.

No. 9913. *Calochortus luteus* Dougl.
Red-brown spot in center
of petal. North Fork Calaveras River
Bridge, n. of San Andreas. Dry gravelly
soil, in dry grass. Gentle open
hillslopes, high with dry grass, are
yellow with this species. It is
very abundant throughout the
region everywhere.

No. 9914. *Delphinium variegatum* T. & G.
With 9913. det R.B. Nov. 59

No. 9915. *Hypochaeris glabra*
Very abundant over the
open hillslopes

[Cont. no. 9960. *Habenaria Calaveras* Co.
Spur twice as long as ovary! Spike
with discrete fls., alternate at apex!
Lip triangular, ^{long-}obtus at tip, minutely ^{*}dentic-
ulate on side! fls. greenish!

* This is very obscure and not of much
moment.

[Cont. 9954, from p. 171. The anthers
have the filaments attached on the
back and prolonged upward into two
slender appendages. These lie close
along the anthers or rather in the
channel between the anther lobes
and are easily overlooked. See p. 177.

Siphocampylodes
No. 9916. *Scutellaria angustifolia*
Pursh. North Fork Calaveras River, near
San Andreas. Lower pair filaments with
the inner edge ciliate; upper pair with
both edges ciliate. All the filaments
are flattened above and all lie
under the upper lip, four side by
side. "Inner" therefore refers to the


edge towards ~~from~~ its pair. The style is glabrous. Upper lip keeled, the lateral lobes reflexed or recurved rather. Corolla glandular-puberulent on the tube outside, with short gland-tipped hairs. The upper stamens have their filaments united with the corolla tube, their edges hairy. The style is quite free and yet it is hidden behind the edges of these stamens and only appears above. See no. 10,002.

[Cont. from p. 171. *Brodiaea*.] only *Chlorogalum pomeridianum*. Cf. also no. 9981, Napa Valley, which is identical ^{of} species and similarly sheathed.

- *Corium maculatum*
 - *Anagallis arvensis*
 - *Melilotus indica*
- Valley Springs.

San Andreas
"

- No. 9917. *Plantago lanceolata*
- No. 9918. *Eschscholzia californica*
Sham. fls. yellow, not golden except in center.
- No. 9919. *Cotyledon lanceolata* B. & W.
var. *incerta* Jepson.
- No. 9920. *Galium*
Abundant on the pastured hills.
- No. 9921. *Verbascum blattaria* L.
Stream bed of the Calaveritas
- No. 9922. *Calochortus albus*
San Andreas, May 26
- No. 9923. *Hypochaeris glabra*
San Andreas, May 26.
- No. 9924. *Filago*
San Andreas, May 26.




— Cont. from p. 193. No. 9961. Plants 3 to 10 ft. broad, 0.8 to 2.4 ft. high. Stems decumbent or semi-prostrate, taking root, and so spreading on all sides from the central stem and thus really becoming a vegetative colony. The main stems are mostly flattened vertically and dead above, the living part being below.

May 27, 1923

[Cont. from p. 175. No. 9919. Corolla livid-crimson, its segments approximate and the corolla therefore acute. I did not see the segments spreading. Filaments filiform. Racemes of the panicle secund, Cotyledon

[Cont. from p. 191. Brodiaea.

See p. 173 also.
" " 171 "



No. 9925. *Pogogyne ziziphorioides* Bth.

Cor. purple; lower lip: middle lobe quadratish, the larger, lateral lobes oblong-quadratish; spreading. Stamens 2, on lower side; ^{all} style ciliate above, staminal fil. inserted very high, the fil. short. Calyx teeth subulate, unequal, bristly ciliate; two teeth long, 3 short. ^{fil.} Neither stamens nor style exerted. Bracts 3-divided into linear or oblanceolate segments, the segments bristly-ciliate. Cor. 4 lines long, violet-purple, a dark purple spot on middle lobe lower lip. Style a little flattened, a little ciliate at apex. Not a trace of upper pair of stamens or anthers. Are there ever?

Calyx teeth as often subequal; commonly nearly as long as cor.; often a little exceeding cor. Lower lip of cor. sometimes showing a little white marking as well as purple. Cor. tube pubescent outside with short spreading hairs.

c. 500 ft. May 28, 1923.

Stigma lobes very unequal. Lower lip with white and purple markings occur as well as just with purple. Plants erect! scarcely "stunish". Scent very sweet. Odor very lovely; better say delightful.

No. 9926. *Pogogyne sarpyllorides* Gray?

Exactly like no. 9925 in most all particulars. More branched, less distinctly erect, more odorous!, calyx-lobes lanceolate, whereas in # 9925 subulate! The difference in appearance of these two calyces is very marked, more than might be supposed from the terms used in indicating the difference. Same habitat as 9925.

No. 9927. *Centunculus minimus* L.

The diagnosis in Fl.-W. Mid. Cal. ed. 2 is satisfactory. Calyx-segments mostly 5 with 4-merous corolla and 4 stamens. Sepals linear-acuminate, mucronate-tipped. - Beds of winter pools.

May 28, 1923.

No. 9928. *Hypericum concinnum*
Stamens in a cluster, 18, 23, 24.

3 Fascicles very distinct. Four of the filaments united below, the other joined only at base. Ovary with 3 placentae, three connivent (but only connivent) in center.

No. 9929. *Downingia*

See p. 187. Upper lobe lanceolate, horizontally recurving. Lower lip: a central white spot on the field of pale blue, a quadratish yellow spot next the throat centering on the white; lobes elliptic, somewhat spreading and thus trefoil shaped. Anther-column nearly straight. Anther apex with 2 small (minute) teeth. Sines between lips cut straight down but not deeply. Sines between lobes of upper lip cut almost as deeply. Calyx - lobes ascending. with 9925

No. 9930 *Gnaphalium palustre* Nutt.
with 9927. Cont. p. 181.

No. 9931. *Chlorogalum angustifolium* Kell.
Fls. dull white. Opening only

at sun-down. Apparently only vesper-tine. Clay flat. See p. 187. Also p. 186.

No. 9932. *Gratiola ebracteata* Bth.
with 9927. See p. 188.

No. 9933. ~~*Loxosiphon*~~ *Pilularia americana*
A.Br. (det. R. Tryon Jr., 1958)
with 9927. See p. 188.

No. 9934. *Boriduralia stricta* ^(A. Gray) Greene.
with 9927. See p. 188.

No. 9935. *Ranunculus*
with 9927. See p. 188.

No. 9936. *Centaurium variegatum* Rob.
See p. 187. Stigmas obovate, obtuse.


Anthers linear, cleft at base, the fls. inserted at summit of cleft. Dry flat.

- Cont. from p. 180. No. 9930. Heads with 2 kinds of corollas: ♀ fls. with filiform corollas and scarcely any limb; ♂ fls. with comparatively large cor. and large limb.
- own -


Involucral bracts obtuse, as says the Fl. W. Mid. Cal., but also subacute; mostly denticulate. Heads mostly ♀, the disk whitish, with a central brown spot of ♂ fls. See p. 189.

No. 9937. *Brodiala minor* (Benth.) Wats. (Or *B. coronaria* (Benth.) Engelm. & Or var. *nana* Hoover) With 9931. Perianth violet-

purple; segments linear-oblong longer than the tube. Anthers as if emminent around style, the staminodes (with revolute edges) approximate around these and much exceeding the anthers. Umbels 1 to 5-flowered.



= anther



= staminode; outline only.
Its edges involute.
Slightly apiculate at the truncatish apex.

dry sandy flat.
See also p. 187.

No. 9938. *Gilia* (*Karwinskii*) *viscidula*. (Bth.) Gray. In hard, very hard, baked, gravelly soil. Fls. red-purple. Calyx glandular, not pubescent, certainly not in any obvious way as in 9941, 9945, 9946. Stamens included. Anthers blue. Corolla throat with (Cont. p. 189).

May 28, 1923.

No. 9939. *Forch's nodosa* Gaertn.

In shade of oaks; out in wild places in the foothills. See p. 187.

No. 9940. *Triteleia* ~~*Brodiala*~~ *hyacinthina*

The most abundant liliaceous plant, at this time, in the lower foothills. Miles and miles of slopes were characterized by its white flowers.

No. 9941. *Gilia* (*Karwinskii*) *pubescens* Stand. Cor. blue, its shape similar to that of no. 9946. Its calyx is like no. 9945; its anthers are also white like 9945 and exerted in about the same way. * San Andreas, 5 mi. west. See p. 189.

No. 9942. *Trifolium obtusiflorum* Hook. Stream bed, North Fork Calaveras River, n. from San Andreas.

* 9941 is very like 9945. The two may be identic save for corolla color, tho the throat in 9945 seems different. Calyx in 9941 is glandular; not in 9945 and 9946.

No. 9943. *Minulus (Eumulus) tricolor* Hartw. With 9927. Corolla lips not very unequal, but the lower the larger. Each lobe of lower lip with a purple spot near base, the palate a field of yellow with purple dots.* Corolla crimson,

No. 9944. *Lythrum hyssopifolia* L.

Petals oblong (broadly), pale lavender**. Fls. $3\frac{1}{2}$ lines long. Stamens***, inserted low on calyx-tube. Stigma capitate. All parts glabrous. With 9927.

* Lobes upper lip with large purple dot at base. Stigmas large, flabelliform. Two lower calyx lobes small. Next pair larger. Upper quite large.


** Better pink-lavender.

*** Stamens 3, 3, 3, 3, 4.

No. 9945. *Gilia* (~~*harvestia*~~ ^{*= N. cotulaefolia*} ~~*harvestia*~~ *harvestia* Eastw.) Corolla blue with white throat & base of lobes, and a prominent purplish spot in throat below base of each lobe. The leaves are pinnatifid incised into linear or filiform divisions, stiff but not truly spinose, tho' the bracts have spinose teeth. San Andreas, 5 mi. west. See p. 187.

Diagram of corolla

See no. 9946.




No. 9946. *Gilia* (~~*harvestia*~~ ^{*harvestia*} ~~*harvestia*~~ *harvestia* Eastw.)

Cor. almost white or pale lavender! Cor. tubular-funnelform; stamens and style exserted. Bracts pinnatifid into rigid elongated filiform divisions which are very prominently "exserted" (that is extended beyond the head). Bracts spinose-pinnate-

lid, densely white-pubescent with straight white hairs on the lower side, that is on the undivided basal part of the bract. Calyx-lobes white-pubescent at base inside, and a little outside. - San Andreas, 5 mi. west.
See p. 187.

Diagram of corolla:
compare with
diagram of no
9945. The two are
not to the same
scale. Stamens
exserted further
in 9946 than in
9945.



Anthers yellow in 9946.
" white in 9945.

- *Chlorogalum angustifolium* Kell.
No style to ovary! See no. 9931.

- This afternoon we drove west. On the dry ^{from San Andreas} thinly wooded hills I collected 9941, 9945, 9946, 9939. We passed Valley Springs; in very hard gravelly soil almost destitute of other vegetation no. 9938 covered the ground with a red-purple flush. We continue west. In a low flat ^{Burson} near ^{San Andreas} *Brodiaea* (no. 9937) *coronaria* (Salish) Engelm. ^(or *B. minor* var. *nana*) det. Hoover. was found; it was abundant in the shade of *Quercus douglasii*. With it was *Centaureum venustum*, no. 9936, and also *Chlorogalum* (no. 9931), this not in ^{open} flower. (On our return near sunset it was ^{in anthesis} quite conspicuous in this dry, sun-scorched spot.) We drove west ^{beyond Burson}. After a few miles I stopped to gather *Gaaphalium palustre*, no. 9930. In the waterless bed of a winter rivulet I saw a stray flower, a vestige, of *Donningia* (no. 9929). In looking

further I found *Gratiola ebracteata* (no. 9932), and then *Minimus* (no. 9943).

Then I chanced on *Ranunculus* (no. 9935) and *Boissiduralis stricta* (no. 9934), all in the bed of this narrow ^{now dry} water way or on its margin. In each case I found at first poor material - but, soon, by looking further better material in ^{former} pool beds. While thus searching, being of course close down to the ground, often on my knees, or hands and knees, I discovered the tiny *Centunculus minimus* (no. 9927), at first - a very little of it, later more abundantly.

Then I saw some curious plants like a minute *Juncus* that has been ringed or slight-stripped and it proved to be *Isotria* (no. 9933).

- All these plants are minute


May 28, 1923.

or small, and all very interesting.

- Cont. from p. 182. [No. 9938.] about 3 lanceolate patterns below each lobe, the color a very light or pale or whitish purple.

Bract.

This type of bract is very different from those in nos. 9941, 9945, 9946.



Entire plant glandular - puberulent
Ovules 2 or 3 in each cell!

See also p. 187.

✓ No. 9947. *Verbascum blattaria* L.
Fls. white. 3-4 ft. tall.

Burson.

✓ No. 9948. *Galium* (*Nanumetia*) *intertexta*
Stend. Cor. pale lavender.

Burson, in moist stream

bed.

No. 9949. *Boissieria densiflora* (Kinds.) S. Wats.

Burson, in moist stream

bed. Plants 33 in. h., the more
robust ones to branching freely at
base. Cor. crimson.

No. 9950. ~~*Brodiaea*~~ ^{*Intelcia*} *hyacinthina*

I see no difference
in fls. between this and the
one of the upland slopes. This grew
in moist bottom near stream and
is much more showy and robust,
the corolla larger.

No. 9951. *Galium*

The stem forked at

base into two main branches. One
of these, equal in length to the
other was cut off near the base
and shows in the stem. Stems
about 4 ft. long. ↑ the base

✓ No. 9951a. *Limnanthus striata*
Jipson. Cor. white with yellowish
center, the petals markedly brown-
purple veined at base. Moist-
stream bed.

No. 9952.

Stream bottom.

✓ No. 9953. *Dicentra chrysantha*
Broken ground; 6½ ft. h.

✓ No. 9954. *Brodiaea appendiculata* Hoover
det. R. F. Hoover
Flat, in very ground near

Burson. It might have been passed
by as *B. coronaria*, which is very
common and making a fine
show all over the country
hereabouts, but it had all but
passed blooming, whereas *B. cor.*

onaria is just coming into full flower.
B. cormaria has erect or nearly
 erect pedicels, this has ^astrongly
 umbellate infl. with the pedicels
 divaricate-curling from the sum-
 mit of the peduncle. Stamens
 much longer than anthers,
 fleshy, incurved edges, the edges
 denticulate! Per. segments elliptic,
 conspicuously broader than in
B. cormaria. This is very similar
 to ^(*B. appendiculata* Hoover) no. 9937. It is the same as
 no. 9957 (^{*B. appendiculata* n. of} ~~Cananche~~) and no.
 9965 (Buena Vista). ^{in Amador Co.} Cont. p. 173, 177, 171, 199.

No. 9955. *Helianthemum scoparium*.
 Nutt. var. ^{vulgare Jepson.} A low hill-top at Burson
 carried a good deal of it. Large
 plants, very fleshy, 4 ft. broad, 14
 to 20 in. high, the stems very
 slender and numerous, making a
 very dense little bush, with
 easily 200,000 flowers and buds in

case of the larger plant.

For geological
 formation see
 vol. 40, p. 12

No. 9956. *Hypericum concinnum* Bth.
 with 9955

No. 9957. *Brodiaea appendiculata* Hoover
 det. Hoover.

At the road forks, n. of
 (north of Cananche) see p. 192 this site is now flooded
 from Mokelumne River bridge. <sup>by Cananche
 in the
 by the
 by the</sup>

cles spreading almost horizontally and
 then curving up, thus very broad; as much as
 6 in. long. Amador Co.

No. 9958. *Arctostaphylos viscida*
 Parry. Buena Vista. Rock-gravelly
 hill-tops, common; 6-8 ft. high.

No. 9959. *Arctostaphylos man-
 zanita* Parry. Buena Vista, Ama-
 dor Co.

No. 9960. *Habenaria elegans*
 See # 173. ^{171 and} Under *Quercus douglasii*
 with no. 9959. H. A.

No. 9960a. *Centaurium venustum* R. & P.
 Buena Vista, Amador Co.

No. 9961. *Arctostaphylos myrti-
 folia* Parry. Buena Vista, Amador Co.
 For geology see above. See vol. 40, p. 1, also.

No. 9962. *Potentilla wilderana* (Parrish)
 = *P. parryi* Greene!!! Stamens 10; filaments dilated, those opp. the petals with narrower filaments. Petals white, obovate, obtuse, toothed or notched laterally at apex, or entire.

No. 9963. *Heterotheca grandiflora* Nutt. *Mokelumne* River bridge near Comanche, but in Amador Co.

No. 9964. *Chlorogalum angustifolium* Kell. Comanche, Calaveras Co.

No. 9965. *Brodiaea*
 Buena Vista.

No. 9966. *Vitis californica* Bth.
 Lone Valley, Amador Co.

No. 9967. *Adenostoma fasciculatum* H. & A. This shrub makes a fine show all thru the foothills at this time. It looks very clean and very attractive; and its flowering panicles are very large. Why should it not be used as a

May 30, 1923

garden shrub? Some shrubs persisted or were spontaneous along a fence line by the roadway between Mokelumne River bridge and Buena Vista. They looked so well-grown and ^{so} good-looking that Miss Saunders thought they must have been planted. But I think not. They were not regularly spaced and no rancher in the foothills would plant this shrub for ornament on a remote portion of his ranch or elsewhere.

Lagoon Valley, June 2, 1923.

No. 9968. *Aristolochia californica* Solano Co.

Climbing a young *Quercus douglasii* H. & A., which is not, I judge, over 15 years old. Fruiting heavily.

- No. 9969. *Monardella villosa*
 Bth., ^{var. ~~torreyana~~ (Eastw.) subseriata} Calistoga, dry wash bank.
 Cor. 7 lines long. Lower lobes diverging,
 all alike, linear; upper lip linear-
 oblong, cleft nearly but not quite
 to the middle, the lobes
 parallel. Stamens equal, exserted;
 filaments glabrous except a little
 hairy at very base. Fil. lavender.
 Cor. tube with downwardly turning
 or somewhat recurving hairs,
 both inside and out.

- No. 9970. *Pentstemon heterophyllus*
 Lindl. Filaments all and style glabrous.
 Cor. 1 in. long, lower lip with 2
 "palates", one opp. each sinus of the
 lower lip. Fertile stamens inserted on very
 base of tube. Upper lobes finally recurving;
 lower lip spreading at right angles to tube.
 Calistoga
 No. 9971. *Clarkia unguiculata*
~~argens~~ Dougl.
 Petals pink, spreading in

pairs right and left. Anthers intense
 terra-cotta; small anthers ^{↑ Large} white. Calistoga.

- No. 9972. *Cytisus monopetalus*
 L. On broken ground at side
 of grade, east-side road, a little s.
 of Calistoga.

- No. 9973. *Holodiscus discolor* Maxim.
 S. of Calistoga, east side road.

- No. 9974. *Lasthenia glaberrima*

This composite grew in a
 marshy spot, s. of Calistoga, east
 side road. ^{↑ in the valley,}

- No. 9975. *Brunella vulgaris* L.

Valley flat near base of hills,
 e. from Yountville. Cont. p. 199.

- No. 9976. *Agoseris ^{grandiflora} plebeia* Greene.
 Fls. heads narrow. Yountville (e.).

- No. 9977. *Beckmannia syzigachne* (Steud.) Fern.
 (Det. Beecher Crumpton, 1922)


Yountville (e.). This grass grew in shallows near

- No. 9978. *Brodiaea laxa* Bth.
^{= Tritelia}

blue form, small flowered. It is
 difficult to say why this is dif-

ferent from the usual form. And yet each time I see it, I feel that there is a varietal difference. It is (a) smaller flowered (perianth. 13 lines long). It is usually a more intense blue - but not dark blue. It tends to grow in semi-shady spots. Anthers azure blue. The style reaches just as far as the base of the segments! The nerves, running down inside tube, from middle of segments, are abruptly and distinctly winged from a point about 4 lines above the base. Unlabeled often with numerous fls.

Corolla very large!



base of perianth laid open, showing the wings on base of veins.

Low hills E. of Yountville.

No. 9979. *Orthocarpus castilleioides*
Oth. Sacs of lower lip as broad as high. Teeth of lower lip, whitish, with a marked purple spot each, towards base, nearly as long as galea. Sae very hairy on inside on ridges of folds. Filaments and style glabrous. Yountville, sr.

No. 9980. *Tragopogon porrifolius*
Napa. Common by roadsides north.

[Cont. no. 9975. Stamens, corolla, style glabrous. All the anthers with a cusp above; the four stamens uniform in this particular].

No. 9981. *Brodiaea appendiculata* Hoover
det. Hoover.

This is the same as no.

9979, collected in same spot. When I visited this little swale not (over)

No. 9982 *Githopsis*

San Andreas, June 3, 1923.
Joyce M. Saunders.

bar. from the mouth of Conn Creek
 I found only one plant just ^{ap. 29}
 coming into flower. The two
 expanded fls. I took (no. 9899).
 On my return to the spot June
 3, I found difficulty in finding
 any plants at all, even the plant
 from which I had clipped the
 two flowers. All about were quantities
 of *Brodiaea* ^{*elegans*} ~~*coronaria*~~ in full
 flower - in full anthesis. Finally
 I found near the same spot
 in the dry grass three plants
 only - no more! It is evidently
 rare. It is exactly the

same as the strange *Brodiaea*
 (*B. appendiculata* Hoover)
 no. 9954 from Calaveras Co. which
 had also passed flowering, where
 in company with *B. elegans*.
 Its anthesis lobes or teeth at top are
 very long. It has two or three
 stems from a bulb. The bulbs are
 deeply seated (9 inches!).

- Chas. Nielsen, Piercy P.O., S. Fork Eel.

- Cont. from p. 145. "never repaid". If
 this whole account was from Mrs. B.,
 then one may qualify it a bit:

- Re. note at bottom this page. The
 three counties Eldorado, Snodgrass and
 Calaveras were in one judicial district
 in 1864. The right name, probably,
 was Three Counties Court House?

- Re. the note in brackets at bottom
 of p. 167. It may be noted
 that the judicial districts
 varied greatly in size and
 in number of counties included.
 Calaveras may have been in a
 4-county district earlier or later.

- Mrs. J. G. Linnon died ^{in Stockton} Jan. 15, 1923. The funeral services were held in Oakland, Jan. 17, 1923. cf. Oakland Tribune, Jan. 16, 1923, p. 10, col. 3.
- Cont. from p. 143. Some two weeks before his arrival Skottsberg wrote to Satchell saying he wished to lecture before the California Botanical Society. As Satchell had nothing to do with the Society, being a nominal member who never attended a meeting, he was obliged to ask me if I would arrange a meeting ^{of the Society} for Dr. Skottsberg to speak, - which I did.

204.

Addresses

- B. B. Davis, Union Lumber Co., Ft. Bragg
- E. Krenger, " " " "
- H. P. Miller, Pacific Lumber Co.
- G. E. Thompson, Scotia " "
- J. Holmes, Eureka Lumber Co.

RICHMOND SAN RAFAEL

FERRY

TIME



TABLE

SUMMER SCHEDULE

.Subject to Change Without Notice.

DAILY

Lv. Richmond	Lv. San Quentin
7:00 a.m.	7:45 a.m.
8:30 a.m.	9:15 a.m.
10:00 a.m.	10:45 a.m.
11:30 a.m.	12:15 p.m.
1:00 p.m.	1:45 p.m.
2:30 p.m.	3:15 p.m.
4:00 p.m.	4:45 p.m.
5:30 p.m.	6:15 p.m.
7:00 p.m.	7:45 p.m.
8:30 p.m.	9:15 p.m.
*10:00 p.m.	*10:45 p.m.

*Sundays and Holidays only

Unsurpassed Grill**Additional Service**

During periods of heavy traffic on Saturdays, Sundays and Holidays and days preceding holidays two boats will be in operation on a

45 MINUTE SERVICE

**RICHMOND SAN RAFAEL
FERRY & TRANSPORTATION CO.**

Phones

Rich. 231, San Rafael 827-W, Mkt 580